

This Provisional PDF corresponds to the article as it appeared upon acceptance. Fully formatted PDF and full text (HTML) versions will be made available soon.

## **TRAIL receptor I (DR4) polymorphisms C626G und A638C are associated with an increased risk for hepatocellular carcinoma (HCC) in HCV-infected patients**

*BMC Cancer* 2012, **12**:85 doi:10.1186/1471-2407-12-85

Christian Korner (christian.koerner@ukb.uni-bonn.de)  
Katarina Riesner (kriesner@uni-bonn.de)  
Benjamin Kramer (benjamin.kraemer@ukb.uni-bonn.de)  
Marianne Eisenhardt (marianne.eisenhardt@ukb.uni-bonn.de)  
Andreas Glassner (andreas.glaessner@ukb.uni-bonn.de)  
Franziska Wolter (fwolter@uni-bonn.de)  
Thomas Berg (thomas.berg@medizin.uni-leipzig.de)  
Tobias Muller (tobiashp.mueller@charite.de)  
Tilman Sauerbruch (tilman.sauerbruch@ukb.uni-bonn.de)  
Jacob Nattermann (jacob.nattermann@ukb.uni-bonn.de)  
Ulrich Spengler (ulrich.spengler@ukb.uni-bonn.de)  
Hans Dieter Nischalke (nischalke@ukb.uni-bonn.de)

**ISSN** 1471-2407

**Article type** Research article

**Submission date** 30 August 2011

**Acceptance date** 8 March 2012

**Publication date** 8 March 2012

**Article URL** <http://www.biomedcentral.com/1471-2407/12/85>

Like all articles in BMC journals, this peer-reviewed article was published immediately upon acceptance. It can be downloaded, printed and distributed freely for any purposes (see copyright notice below).

Articles in BMC journals are listed in PubMed and archived at PubMed Central.

For information about publishing your research in BMC journals or any BioMed Central journal, go to

<http://www.biomedcentral.com/info/authors/>

© 2012 Korner *et al.*; licensee BioMed Central Ltd.

This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

# TRAIL receptor I (*DR4*) polymorphisms C626G und A638C are associated with an increased risk for hepatocellular carcinoma (HCC) in HCV-infected patients

Christian Körner,<sup>Aff1</sup>

**Email:** christian.coerner@ukb.uni-bonn.de

Katarina Riesner,<sup>Aff1</sup>

**Email:** kriesner@uni-bonn.de

Benjamin Krämer,<sup>Aff1</sup>

**Email:** benjamin.kraemer@ukb.uni-bonn.de

Marianne Eisenhardt,<sup>Aff1</sup>

**Email:** marianne.eisenhardt@ukb.uni-bonn.de

Andreas Glässner,<sup>Aff1</sup>

**Email:** andreas.glaessner@ukb.uni-bonn.de

Franziska Wolter,<sup>Aff1</sup>

**Email:** fwolter@uni-bonn.de

Thomas Berg,<sup>Aff2</sup>

**Email:** thomas.berg@medizin.uni-leipzig.de

Tobias Müller,<sup>Aff3</sup>

**Email:** tobiashp.mueller@charite.de

Tilman Sauerbruch,<sup>Aff1</sup>

**Email:** tilman.sauerbruch@ukb.uni-bonn.de

Jacob Nattermann,<sup>Aff1</sup>

**Email:** jacob.nattermann@ukb.uni-bonn.de

Ulrich Spengler,<sup>Aff1</sup>

**Email:** ulrich.spengler@ukb.uni-bonn.de

Hans Dieter Nischalke,<sup>Aff1</sup>

Corresponding Affiliation: Aff1

**Phone:** +49-228-28751416

**Fax:** +49-228-28751419

**Email:** nischalke@ukb.uni-bonn.de

Aff1 Department of Internal Medicine I, University of Bonn, Sigmund-Freud-Str. 25, 53127 Bonn, Germany

Aff2 Department of Gastroenterology, University Hospital Leipzig, Philipp-Rosenthal-Strasse 27, 04103 Leipzig, Germany

Aff3 Medical Clinic for Hepatology and Gastroenterology, Medical

## Abstract

### Background

Tumour surveillance via induction of TRAIL-mediated apoptosis is a key mechanism, how the immune system prevents malignancy. To determine if gene variants in the TRAIL receptor I (*DR4*) gene affect the risk of hepatitis C virus-(HCV)-induced liver cancer (HCC), we analysed *DR4* mutations C626G (rs20575) and A683C (rs20576) in HCV-infected patients with and without HCC.

### Methods

Frequencies of *DR4* gene polymorphisms were determined by LightSNiP assays in 159 and 234 HCV-infected patients with HCC and without HCC, respectively. 359 healthy controls served as reference population.

### Results

Distribution of C626G and A683C genotypes were not significantly different between healthy controls and HCV-positive patients without HCC. *DR4* variants 626 C and 683A occurred at increased frequencies in patients with HCC. The risk of HCC was linked to carriage of the 626 C allele and the homozygous 683AA genotype, and the simultaneous presence of the two risk variants was confirmed as independent HCC risk factor by Cox regression analysis (Odds ratio 1.975, 95% CI 1.205-3.236;  $p=0.007$ ). Furthermore HCV viral loads were significantly increased in patients who simultaneously carried both genetic risk factors ( $2.69 \pm 0.36 \times 10^6$  IU/ml vs.  $1.81 \pm 0.23 \times 10^6$  IU/ml,  $p=0.049$ ).

### Conclusions

The increased prevalence of patients with a 626 C allele and the homozygous 683 AA genotype in HCV-infected patients with HCC suggests that these genetic variants are a risk factor for HCC in chronic hepatitis C.

## Keywords

TRAIL receptor I, DR4, Apoptosis, Polymorphism, C626G (rs20575), A683C (rs20576), HCV, HCC, Cancer

# Background

Nearly 3% of the world population, app. 180 million people, suffer from hepatitis C virus infection (HCV) thus representing a global health problem [1]. In most of the cases exposure to HCV results in chronic viral persistence. Progression of chronic hepatitis C leads to liver fibrosis and cirrhosis and is associated with an increased risk to develop hepatocellular carcinoma (HCC) [2,3]. HCC has become the third leading cause of cancer-related death worldwide [4–6], and in Western Countries chronic hepatitis accounts for the majority of HCCs [Lit]. HCV proteins can interact with tumour suppressor proteins as well as with proteins involved in cell-cycle control, and these interactions may promote the development of abnormal cells in the liver.

Tumour development is normally prevented by the immune system, which eliminates transformed cells via induction of apoptosis by tumour necrosis factor-related apoptosis-inducing ligand (TRAIL) [7,8]. Binding of TRAIL to its cognate death receptors DR4 and DR5 triggers activation of the apoptotic cascade, formation of apoptotic bodies and eventually to depletion of the apoptotic cell. DR4 and DR5 are members of the tumour necrosis factor super family and are characterized by the existence of an extra-cellular cysteine-rich binding domain as well as an intra-cellular death domain essential for the transmission of the apoptotic stimulus.

Genetic alterations in death receptors might compromise apoptotic cell signalling and therefore contribute to the development of tumour cells. Several studies suggested an increased risk for cancer associated with single nucleotide polymorphisms (SNPs) in the *DR4* gene [9–13]. However, the potential influence of *DR4* gene mutations on the development of HCC has not been investigated so far. Therefore, we analysed the effect of the *DR4* polymorphisms C626G (Thr209Arg, rs20575) and A638C (Glu228Ala, rs20576) on the occurrence of HCC in patients chronically infected with HCV.

# Methods

## Design and study population

A total of 393 patients with chronic hepatitis C virus genotype 1 infection from the Bonn and Berlin University Departments of Gastroenterology were enrolled into this study including 159 patients with HCV-associated HCC and 234 patients without HCC. 359 HCV-negative healthy individuals and 56 patients with HCC caused by chronic hepatitis B served as control groups. All subjects in this study were of Caucasian descent. Further clinical and demographic characteristics are listed in Table 1.

**Table 1** Patient characteristics and distribution of DR4 genotypes

Healthy controls	HCV(+) all without HCC	HCV(+) Cirrhosis (-) without HCC	HCV(+) Cirrhosis (+) without HCC	HCV(+) Cirrhosis (+) with HCC	HBV(+) With HCC
------------------	------------------------	----------------------------------	----------------------------------	-------------------------------	-----------------

<b>Numbers</b>	359	234	159	75	159	56
<b>Median age (range)</b>	59 (35-69)	52 (39-63)	46 (39-57)	55 (49-63)	62 (54-70)	60(24-83)
<b>Male Gender (%)</b>	51.4	66.2	65.2	69.7	58.6	75.0
<b>HCV infection (%)</b>	-	100	100	100	100	0
<b>HCV genotype 1 (%)</b>	-	100	100	100	100	-
<b>HCV viral load (10<sup>6</sup>IU/ml)<sup>a)</sup></b>	-	2.04 ± 0.25	2.10 ± 0.29	1.86 ± 0.48	1.88 ± 0.36	-
<b>HBV infection (%)</b>	-	3.2	2.8	4.0	4.4	100
<b>BMI &gt; 30 (%)</b>	-	17.1	15.0	21.7	26.2	21.4
<b>Diabetes (%)</b>	-	7.7	0.8	23.0	12.0	3.6
<b>rs20575 (C626G)<sup>b)</sup></b>	349 (100)	233 (100)	159 (100)	74 (100)	159 (100)	56 (100)
<b>CC</b>	62 (17.8)	56 (24.0)	38 (23.9)	18 (24.3)	40 (25.2)	9 (16.1)
<b>CG</b>	166 (47.6)	108 (46.4)	74 (46.5)	34 (45.9)	87 (54.7)	30 (53.6)
<b>GG</b>	121 (34.7)	69 (29.6)	47 (29.6)	22 (29.7)	32 (20.1)	17 (30.4)
<b>Deviation from Hardy-Weinberg equilibrium<sup>c)</sup></b>	P=0.74	P=0.29	P=0.43	P=0.49	P=0.27	P=0.59
<b>rs20576 (A683C)<sup>b)</sup></b>	359 (100)	234 (100)	159 (100)	75 (100)	159 (100)	56 (100)
<b>CC</b>	19 (5.3)	10 (4.3)	6 (3.8)	4 (5.3)	5 (3.1)	1 (1.8)
<b>AC</b>	125 (34.8)	85 (36.3)	59 (35.0)	26 (34.7)	33 (21.9)	17 (30.4)
<b>AA</b>	215 (59.9)	139 (59.4)	94 (59.1)	45 (60.0)	121 (75.8)	38 (67.9)
<b>Deviation from Hardy-Weinberg equilibrium<sup>c)</sup></b>	P=0.88	P=0.58	P=0.49	P=1.0	P=0.17	P=1.0

<sup>a)</sup> Mean ± SEM, <sup>b)</sup> Number of patients (number/ % of total), <sup>c)</sup> Fisher's exact test

Cirrhosis was diagnosed either by liver biopsy, transient elastography (stiffness >15 kPa), and signs of portal hypertension (splenomegaly, esophageal varices, ascites). The diagnosis of HCC was made by contrast enhanced magnetic resonance imaging and computed tomography according to recently established diagnostic criteria [EASL 2009 und AASLD 2010 guidelines].

Informed consent was obtained from all patients. The study conformed to the ethical guidelines of the Helsinki declaration and had been approved by the University of Bonn ethics committee (reference number: 019/07).

## Diagnosis of HCV Infection

HCV antibodies were detected with a micro particle enzyme immunoassay (AxSYM; Abbott) and confirmed by dot immunoassay (Matrix; Abbott). HCV RNA was detected with a nucleic acid purification kit (QIAamp Viral Kit; Qiagen, Hilden, Germany), followed by reverse transcription and nested polymerase chain reaction. Quantitative determination of HCV loads was done by branched DNA technology (Chiron, Emeryville, CA). HCV genotype was determined by the Innolipa II line probe assay (Innogenetics, Zwijndrecht, Belgium).

## DR4 Genotyping

Genomic DNA was extracted from 200 µl EDTA-blood using the QIAamp Blood Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's protocol. Determination of the *DR4* gene polymorphisms C626G (rs20575) and A683C (rs20576) were performed by LightCycler real time PCR (Roche, Mannheim, Germany) using commercial LightSNiP (SimpleProbe) assays from TIB-MolBiol (Berlin, Germany) according to the manufacturer's recommendations.

## Statistical analysis

Genotype frequencies were determined and tested for consistency with Hardy-Weinberg equilibrium using an exact test. Allele and genotype frequencies were compared between cases and controls by Pearson's goodness-of-fit ( $\chi^2$ ) test and Armitage's trend test, respectively (<http://ihg2.helmholtz-muenchen.de/cgi-bin/hw/hwa1.pl>).

Statistical analysis was performed with SPSS 18.0 (SPSS, Munich, Germany). Data are given as means  $\pm$  SD, unless stated otherwise. Differences between groups were analyzed by t-test and Mann-Whitney-U test as appropriate.

To take into account potentially confounding risk factors of cirrhosis (age, gender, HBV infection, alcohol and obesity), univariate comparisons (ANOVA and chi<sup>2</sup>-statistics) followed by forward conditional logistic regression were performed. Parameters with univariate effects at  $p < 0.1$  were entered into the multivariate analysis with  $P < 0.05$  for inclusion and  $p > 0.1$  for exclusion as selection criterion for parameters in the final statistical model.

## Results

The distribution of DR4 C626 > G and A683 > C genetic variants matched the Hardy-Weinberg equilibrium in all analysed groups and subgroups. HCV-infected patients without HCC revealed a slightly higher allele frequency of the DR4 626 C allele (47.2%)-corresponding to an increased prevalence of 626 C carriers (70.4%)-than our healthy controls (allele frequency 41.5%; frequency of 626 C carriers 65.3%; Figure 1A). These differences within the HCV-infected patient groups were apparently independent from the presence of cirrhosis and did not reach statistical significance.

### **Figure 1 Frequencies of DR4 risk alleles in HCV-infected patients with and without HCC.**

Figure A shows the frequency of carriers with a *DR4* 626 C risk allele, and figure B the frequency of patients with the homozygous *DR4* 683AA genotype. Figure C shows the frequency of patients who carried either the 626CG or 626CC risk genotype in combination with the homozygous *DR4* 683AA genotype. Differences between the various patient groups were analyzed by chi<sup>2</sup> statistics.

Prevalence of carriers with a 626 C allele (genotypes 626CG and CC) (79.9%) and 626 C allele frequency (52.5%) were significantly increased in the patients with HCV-associated HCC (Figure 1A). The 626 C allele frequency in patients with HCV-associated HCC was significantly different only from that in healthy controls (OR = 1.556, 95%CI: 1.192-2.031,  $p = 0.001$ ).

Importantly, the increased prevalence of carriers with a 626 C allele reached statistical significance both with respect to healthy controls (OR = 2.106, 95%CI: 1.348-3.290,  $p = 0.001$ ) and to HCV-infected patients without HCC (OR = 1.670, 95%CI: 1.034-2.696,  $p = 0.034$ ) (Figure 1A).

The distribution of A683 > C variants did not differ between healthy controls and HCV-infected patients without HCC irrespective from the presence of cirrhosis (Table 1). Patients with HCV-associated HCC had the highest frequency of the 623A allele (86.5%), which was significantly increased as compared to healthy controls (A allele frequency 77.3%, OR = 1.856 95% CI: 1.291-2.667,  $p = 0.0006$ ) and HCV-infected patients without HCC (A allele frequency 77.6%, OR = 1.828, 95%CI: 1.243-2.686,  $p = 0.002$ ). This deviation was strong enough to achieve statistical significance even when HCV-infected subgroups with and without cirrhosis were compared separately (cirrhosis: 77.3% A allele frequency, OR = 1.852, 95%CI: 1.126-3.045,  $p = 0.013$ ; without cirrhosis: 77.7% A allele frequency, OR = 1.816, 95%CI: 1.201-2.746,  $p = 0.004$ ).

These changes were due to a significant increase in the proportion of patients with the homozygous 683AA genotype in patients with HCV associated HCC (76.1%) as compared to healthy controls (59.9%, OR = 0.477, 95%CI: 0.314-0.725,  $p = 0.0003$ ) and HCV-infected patients without HCC (59.4%, OR = 0.468, 95% CI: 0.300-0.730,  $p = 0.0006$ ), irrespective whether cirrhosis was present (60.0%, OR = 0.480, 95%CI: 0.267-0.862,  $p = 0.011$ ) or not (59.1%, OR = 0.462, 95%CI: 0.286-0.747,  $p = 0.001$ ) (Figure 1B). Interestingly, HCV viral loads tended to be increased in patients who carried the genetic risk variants associated with polymorphisms C626G ( $2.44 \pm 0.28 \times 10^6$  IU/ml vs.  $1.85 \pm 0.30 \times 10^6$  IU/ml, n.s.) and A683C ( $2.30 \pm 0.27 \times 10^6$  IU/ml vs.  $1.94 \pm 0.26 \times 10^6$  IU/ml, n.s.), and this effect reached statistical significance in patients who simultaneously carried both genetic risk factors ( $2.69 \pm 0.36 \times 10^6$  IU/ml vs.  $1.81 \pm 0.23 \times 10^6$  IU/ml,  $p = 0.049$ ) (Figure 2).

**Figure 2 HCV viral loads in patients with and without genetic DR4 risk variants.** This Figure shows HCV viral loads (Means  $\pm$  SEM) in patients who carry the DR4 626CG or 626CC risk variant (left columns), the DR4 683AA risk variant (middle columns) and both DR4 risk variants simultaneously (right columns). Dark columns refer to patients with the studied risk factor, grey columns to those without the risk factor, respectively. Differences were compared by ANOVA.

Next, we checked whether variants in the TRAIL receptor I (DR4) gene were confirmed as independent HCC risk factors in HCV-infected patients when known risk factors such as age, sex, obesity, diabetes, HCV viral load and HBV co-infection were also taken into account (Table 2). Univariate analysis identified age (OR = 1.076 per year, 95%CI: 1.053-1.098,  $p < 0.0001$ ), carriage of a 626 C allele (OR = 1.670, 95%CI: 1.034-2.696,  $p = 0.035$ ), the homozygous 683AA genotype (OR = 2.176, 95%CI: 1.390-3.407,  $p = 0.001$ ) and the simultaneous presence of both genetic variants (OR = 1.940, 95%CI: 1.280-2.940,  $p = 0.002$ ) as risk factors of HCC (Table 2). In the multivariate regression analysis age was found to be an independent risk factor for the development of HCC in chronic hepatitis C (OR = 1.074, 95%CI: 1.052-1.097,  $p \leq 0.001$ ). More importantly, we could confirm that the simultaneous presence of a 626 C allele in combination with the homozygous 683AA genotype was significantly associated with the occurrence of HCC

in this continuative analysis (OR = 1.975, 95%CI: 1.205-3.236,  $p = 0.007$ ) and could therefore identify the combined carriage of a DR4 risk variant as new independent predictor for the development of HCC in chronic hepatitis C.

**Table 2** Regression analysis for risk factors of HCC among patients with hepatitis C genotype 1

<b>Univariate analysis</b>				
Parameter	p-value	Odds ratio	95 % Confidence interval	
			Lower	Upper
Age §	0.000	1.076	1.053	1.098
Diabetes mellitus	0.369	1.412	0.691	2.886
Gender (male)	0.135	1.430	0.915	2.236
HBV infection	0.626	1.295	0.460	3.646
HCV viral load ( $10^6$ IU/ml)	0.339	0.960	0.883	1.044
Obesity (BMI >30)	0.153	1.668	0.867	3.211
C626G genotype ([CC+CG] vs. GG)	0.035	1.670	1.034	2.696
A683C genotype (AA vs. [AC+CC])	0.001	2.176	1.390	3.407
Combined 626C and 683AA risk genotype vs. all other combinations	0.002	1.940	1.280	2.940
<b>Multivariate analysis*</b>				
Parameter	p-value	Odds ratio	95 % Confidence interval	
			Lower	Upper
Age §	0.000	1.074	1.052	1.097
Combined 626C and 683AA risk genotype	0.007	1.975	1.205	3.236

\* including all significant parameters from univariate analysis, § OR (95% CI) per year

To check, if our findings were specific for hepatitis C, we performed a similar analysis in our group of HCC patients, who had chronic hepatitis B. However, unlike hepatitis C-associated HCC the distributions of genetic DR4 variants in HBV-associated HCC, healthy controls and HCV-positive patients without HCC were not significantly different (Table 1). In particular, the simultaneous combination of a 626 C allele with the 683AA genotype was observed in rather

similar frequencies in the groups with HBV-associated HCC (50.0%), hepatitis C without HCC (49.4%) and healthy controls (47.0%) while the difference in frequency between HBV- and HCV-associated HCC (65.4%) was significant (OR = 1.891, 95%CI: 1.020-3.506, p = 0.042).

## Discussion

Emerging evidence suggest an important role of TRAIL for control and elimination of HCV infection. TRAIL has been implicated in the death of HCV-infected but not normal liver cells [14]. In HCV infection its expression on natural killer cells is up-regulated by interferon, and is inversely correlated to HCV-RNA serum levels [15,16]. TRAIL can also induce cell death in hepatic tumour cells but expression of TRAIL receptors on human hepatocellular carcinoma is variable and frequently down-regulated [17]. Nevertheless, several studies suggest TRAIL receptor-mediated apoptosis to play a role in the elimination of tumour cells in human hepatocellular carcinoma [18,19].

In this study we found that the TRAIL receptor I wild type with threonine at amino acid position 209 (626 C) and alanine at position 228 (638A) is associated with an increased risk of HCC in patients with chronic hepatitis C. Of note, it was particularly the combination of the risk genotypes 626CC or 626CG with the homozygous 683AA genotype that was identified as a new independent predictor for HCC. Interestingly, this association of DR4 genetic variants and the risk to develop HCC was evident only for patients with chronic hepatitis C but not hepatitis B, suggesting a critical role of the aetiology underlying HCC. Both polymorphisms in the *DR4* gene have already been reported to affect the risk for other types of malignancy. However, in a previous study investigating the role of the *DR4* A683C polymorphism in chronic lymphocytic leukaemia, mantle cell lymphoma, prostate cancer, head and neck squamous cell carcinoma and bladder cancer an increased frequency of the CC genotype was apparently linked to malignancy [20]. In contrast, Frank *et al.* observed an increased percentage of carriers with the AA genotype in patients with breast cancer [10]. Moreover, they found an increased percentage of carriers of the 626CC genotype similar to our results. At present, it remains unclear which mechanisms underlie *DR4* polymorphisms C626G and A683C to affect the risk for malignancy.

However, both amino acid exchanges are in the extracellular cysteine-rich domain of DR4. Therefore both genetic variants may lead to alterations in the TRAIL-binding domain and thus alter DR4 affinity for TRAIL. Since TRAIL signalling presumably contributes importantly to the control of HCV infection, our finding of significantly increased HCV viral loads in carriers of both *DR4* risk factors suggests that TRAIL-DR4 signalling is less efficient in these patients. In line with this reasoning less efficient signalling of cell death in transformed cells and reduced susceptibility of transformed hepatocytes towards TRAIL-induced apoptosis would also facilitate HCC development. On the other hand, we cannot exclude the possibility that the increased HCC risk in HCV-infected patients carrying the C626G and A683C risk variants of TRAIL receptor I simply reflects less efficient immune control over HCV infection via TRAIL-mediated mechanisms. TRAIL is likely to affect tumour surveillance of transformed HCV-infected cells by the immune system and viral loads, once HCV infection has occurred [14]. On the other hand our data suggest that neither of the *DR4* polymorphisms affects susceptibility to HCV infection. This constellation can be explained by the fact that DR4 expression is rather low on normal liver cells while up-regulated DR4 expression following HCV infection sensitizes liver cells towards

TRAIL-mediated apoptosis [14,21]. Thus, sufficient DR4 expression on transformed liver cells appears to be a pivotal prerequisite for efficient tumour surveillance by the immune system, in line with recent clinical findings [17].

Therefore the role of TRAIL and *DR4* polymorphisms should be further studied in patients whose risk for HCC is attributed to different aetiologies.

## Conclusion

Here, we provide first evidence that HCV genotype 1 infected carriers of a DR4 626 C allele in combination with the DR4 683AA genotype have an increased risk for HCC indicating that TRAIL and its receptors contribute importantly to control of HCV infection and tumour surveillance by the immune system. The combined presence of the two DR4 risk variants can help to identify HCV-infected patients with an increased risk for liver cancer, who may need more intensive cancer monitoring.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

K.R. and F.W. established and performed the real-time PCRs for DR4 genotyping. B. K., M. E., T. M. and T. B. collected and analyzed clinical data and collected samples from the study population. H.D.N., U.S., and J.N. designed the study and did statistical analysis. C.K. and T.S. wrote the manuscript. All authors read and approved the final manuscript.

## Acknowledgements

This work was supported by a grant from the German Federal Ministry of Education and Research (BMBF) within the network for resistance in hepatitis C, grant number (01KI0792) to JN and US. By a grant from the H.W. & J. Hector foundation, grant number (M42) to JN and by the Deutsche Krebshilfe grant number (107865) to HDN and US.

## References

1. Alter MJ: **Epidemiology of hepatitis C.** *Hepatology* 1997, **26**(3 Suppl 1):62S–65S.
2. Moradpour D, Blum HE: **Pathogenesis of hepatocellular carcinoma.** *Eur J Gastroenterol Hepatol* 2005, **17**(5):477–483.
3. Tsukuma H, Hiyama T, Tanaka S, Nakao M, Yabuuchi T, Kitamura T, Nakanishi K, Fujimoto I, Inoue A, Yamazaki H, *et al*: **Risk factors for hepatocellular carcinoma among patients with chronic liver disease.** *N Engl J Med* 1993, **328**(25):1797–1801.

4. Altekruse SF, McGlynn KA, Reichman ME: **Hepatocellular carcinoma incidence, mortality, and survival trends in the United States from 1975 to 2005.** *J Clin Oncol* 2009, **27**(9):1485–1491.
5. Schutte K, Bornschein J, Malfertheiner P: **Hepatocellular carcinoma—epidemiological trends and risk factors.** *Dig Dis* 2009, **27**(2):80–92.
6. Shepard CW, Finelli L, Alter MJ: **Global epidemiology of hepatitis C virus infection.** *Lancet Infect Dis* 2005, **5**(9):558–567.
7. Mellier G, Huang S, Shenoy K, Pervaiz S: **TRAILing death in cancer.** *Mol Aspects Med* 2010, **31**(1):93–112.
8. Yang A, Wilson NS, Ashkenazi A: **Proapoptotic DR4 and DR5 signaling in cancer cells: toward clinical translation.** *Curr Opin Cell Biol* 2010, **22**(6):837–844.
9. Chen B, Liu S, Wang XL, Xu W, Li Y, Zhao WH, Wu JQ: **TRAIL-R1 polymorphisms and cancer susceptibility: an evidence-based meta-analysis.** *Eur J Cancer* 2009, **45**(14):2598–2605.
10. Frank B, Hemminki K, Shanmugam KS, Meindl A, Klaes R, Schmutzler RK, Wappenschmidt B, Untch M, Bugert P, Bartram CR, *et al*: **Association of death receptor 4 haplotype 626 C-683C with an increased breast cancer risk.** *Carcinogenesis* 2005, **26**(11):1975–1977.
11. Frank B, Shanmugam KS, Beckmann L, Hemminki K, Brenner H, Hoffmeister M, Chang-Claude J, Burwinkel B: **Death receptor 4 variants and colorectal cancer risk.** *Cancer Epidemiol Biomarkers Prev* 2006, **15**(10):2002–2005.
12. Ulybina YM, Kuligina E, Mitiushkina NV, Rozanov ME, Ivantsov AO, Ponomariova DN, Togo AV, Levchenko EV, Shutkin VA, Brenister SI, *et al*: **Coding polymorphisms in Casp5, Casp8 and DR4 genes may play a role in predisposition to lung cancer.** *Cancer Lett* 2009, **278**(2):183–191.
13. Wang M, Wang M, Cheng G, Zhang Z, Fu G, Zhang Z: **Genetic variants in the death receptor 4 gene contribute to susceptibility to bladder cancer.** *Mutat Res* 2009, **661**(1–2):85–92.
14. Lan L, Gorke S, Rau SJ, Zeisel MB, Hildt E, Himmelsbach K, Carvajal-Yepes M, Huber R, Wakita T, Schmitt-Graeff A, *et al*: **Hepatitis C virus infection sensitizes human hepatocytes to TRAIL-induced apoptosis in a caspase 9-dependent manner.** *J Immunol* 2008, **181**(7):4926–4935.
15. Ahlenstiel G, Edlich B, Hogdal LJ, Rotman Y, Nouredin M, Feld JJ, Holz LE, Titerence RH, Liang TJ, Rehermann B: **Early Changes in Natural Killer Cell Function Indicate**

**Virologic Response to Interferon Therapy for Hepatitis C.** *Gastroenterology* 2011, **141**(4):1231–1239.

16. Stegmann KA, Bjorkstrom NK, Veber H, Ciesek S, Riese P, Wiegand J, Hadem J, Suneetha PV, Jaroszewicz J, Wang C *et al*: **Interferon-alpha-induced TRAIL on natural killer cells is associated with control of hepatitis C virus infection.** *Gastroenterology* 2010, **138**(5):1885–1897.

17. Kriegl L, Jung A, Engel J, Jackstadt R, Gerbes AL, Gallmeier E, Reiche JA, Hermeking H, Rizzani A, Bruns CJ *et al*: **Expression, cellular distribution, and prognostic relevance of TRAIL receptors in hepatocellular carcinoma.** *Clin Cancer Res* 2010, **16**(22):5529–5538.

18. Chen XP, He SQ, Wang HP, Zhao YZ, Zhang WG: **Expression of TNF-related apoptosis-inducing Ligand receptors and antitumor tumor effects of TNF-related apoptosis-inducing Ligand in human hepatocellular carcinoma.** *World J Gastroenterol* 2003, **9**(11):2433–2440.

19. He S, Chen Y, Chen X, Zhao Y, Wang H, Zhang W, Wang S: **Antitumor effects of soluble TRAIL in human hepatocellular carcinoma.** *J Huazhong Univ Sci Technolog Med Sci* 2005, **25**(1):51–54.

20. Wolf S, Mertens D, Pscherer A, Schroeter P, Winkler D, Grone HJ, Hofele C, Hemminki K, Kumar R, Steineck G, *et al*: **Ala228 variant of trail receptor 1 affecting the ligand binding site is associated with chronic lymphocytic leukemia, mantle cell lymphoma, prostate cancer, head and neck squamous cell carcinoma and bladder cancer.** *Int J Cancer* 2006, **118**(7):1831–1835.

21. Zhu H, Dong H, Eksioglu E, Hemming A, Cao M, Crawford JM, Nelson DR, Liu C: **Hepatitis C virus triggers apoptosis of a newly developed hepatoma cell line through antiviral defense system.** *Gastroenterology* 2007, **133**(5):1649–1659.

Figure 1




Figure 2

