

This Provisional PDF corresponds to the article as it appeared upon acceptance. Fully formatted PDF and full text (HTML) versions will be made available soon.

PKCalpha expression is a marker for breast cancer aggressiveness

Molecular Cancer 2010, 9:76 doi:10.1186/1476-4598-9-76

Gry Kalstad Lonne (Gry.Kalstad_Lonne@med.lu.se) Lousie Cornmark (Lousie.Cornmark@med.lu.se) Iris Omanovic Zahirovic (iris_zahirovic@yahoo.se) Goran Landberg (Goran.Landberg@med.lu.se) Karin Jirstrom (Karin.Jirstrom@med.lu.se) Christer Larsson (Christer.Larsson@med.lu.se)

ISSN	1476-4598
Article type	Research
Submission date	3 November 2009
Acceptance date	14 April 2010
Publication date	14 April 2010
Article URL	http://www.molecular-cancer.com/content/9/1/76

This peer-reviewed article was published immediately upon acceptance. It can be downloaded, printed and distributed freely for any purposes (see copyright notice below).

Articles in *Molecular Cancer* are listed in PubMed and archived at PubMed Central.

For information about publishing your research in *Molecular Cancer* or any BioMed Central journal, go to

http://www.molecular-cancer.com/info/instructions/

For information about other BioMed Central publications go to

http://www.biomedcentral.com/

© 2010 Kalstad Lonne et al., licensee BioMed Central Ltd.

This is an open access article distributed under the terms of the Creative Commons Attribution License (<u>http://creativecommons.org/licenses/by/2.0</u>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

PKCα expression is a marker for breast cancer aggressiveness

Gry Kalstad Lønne, Louise Cornmark, Iris Omanovic Zahirovic, Göran Landberg, Karin Jirström, and Christer Larsson*

Center for Molecular Pathology, Department of Laboratory Medicine, Lund University, Malmö University Hospital, SE- 205 02 Malmö, Sweden

*Corresponding author: Christer Larsson Center for Molecular Pathology Dept of Laboratory Medicine Lund University Malmö University Hospital Entrance 78, 3rd floor SE- 205 02 Malmö Sweden Tel.: +46-(0)40-337404 Fax.: +46-(0)40-337063 christer.larsson@med.lu.se

Email addresses:

- GKL: gry.kalstad_lonne@med.lu.se
- LC: louise.cornmark@med.lu.se
- IOZ: iris_zahirovic@yahoo.se
- GL: goran.landberg@med.lu.se
- KJ: karin.jirstrom@med.lu.se
- CL: christer.larsson@med.lu.se

Abstract

BACKGROUND: Protein kinase C (PKC) isoforms are potential targets for breast cancer therapy. This study was designed to evaluate which PKC isoforms might be optimal targets for different breast cancer subtypes.

RESULTS: In two cohorts of primary breast cancers, PKC α levels correlated to estrogen and progesterone receptor negativity, tumor grade, and proliferative activity, whereas PKC δ and PKC ϵ did not correlate to clinicopathological parameters. Patients with PKC α positive tumors showed poorer survival than patients with PKC α -negative tumors independently of other factors. Cell line studies demonstrated that PKC α levels are high in MDA-MB-231 and absent in T47D cells which proliferated slower than other cell lines. Furthermore, PKC α silencing reduced proliferation of MDA-MB-231 cells. PKC α inhibition or downregulation also reduced cell migration *in vitro*.

CONCLUSIONS: PKC α is a marker for poor prognosis of breast cancer and correlates to and is important for cell functions associated with breast cancer progression.

Background

Breast cancer is a heterogeneous disease which encompasses several subgroups with different morphology, genetic changes, and response to therapies [1, 2]. It is therefore important to gain more insight into relevant therapeutic targets for each subgroup to optimize tailored treatment protocols for individual patients. Numerous intracellular signaling proteins have been suggested to be promising targets for blocking the malignancy of breast cancer cells. The protein kinase C (PKC) isoforms are examples of such potential therapeutic targets.

PKC is a family of serine/threonine kinases involved in several processes including proliferation, differentiation, apoptosis, and migration. The PKC isoforms are divided into three subgroups depending on the structure of the regulatory domain: classical (PKC α , β I, β II, and γ), novel (PKC δ , ε , and θ), and atypical (PKC ζ and ν/λ) isoforms. Classical and novel PKCs contain a diacylglycerol (DAG)-binding C1 domain and are therefore regulated by activation of pathways that lead to DAG generation. Atypical PKCs are DAG-insensitive and regulated in a different manner [3].

Several studies have implicated the DAG-sensitive classical and novel PKC isoforms in promoting malignant features of breast cancer cells. PKC α has been coupled to estrogen receptor (ER) negativity [4] and estrogen-independent growth of cultured cells [5, 6] and patients with PKC α -negative tumors had better response to endocrine treatment compared to patients with PKC α -positive tumors [7, 8]. Moreover, increased PKC α expression leads to a more aggressive phenotype [4] and is associated with resistance to cytostatic drugs in MCF-7 cells [9, 10]. PKC α is also evaluated as a therapeutic target for breast cancer [11]. However, PKC α levels are reduced in breast

cancer compared to normal breast tissue [12, 13]. Thus, there is evidence for both a promoting and a suppressing role for PKC α in breast cancer.

The role of PKC δ in breast cancer is ambiguous. Patients with PKC δ -positive tumors show better endocrine response compared to patients with PKC δ -negative tumors [8] and PKC δ has been shown to be crucial for UV light-induced apoptosis of cultured breast cancer cells [14]. However, several studies point to a pro-tumorigenic role of PKC δ in breast cancer. PKC δ can induce resistance to tamoxifen and irradiation in cultured breast cancer cells [15, 16] and has been shown to promote both metastasis [17-19] and proliferation [20] of murine mammary cancer and epithelial cells. We have recently shown that depletion of PKC δ is sufficient to drive breast cancer cells into apoptosis [21].

PKCε has frequently been assigned oncogenic effects in breast cancer. Expression levels of PKCε have been shown to correlate with tumor grade, HER2 expression, ER negativity, and poor survival in breast cancer patients. Moreover, in MDA-MB-231 breast cancer cells, downregulation of PKCε reduced the tumor growth and metastatic capacity in mice [22]. There is also evidence that PKCε protects cells against apoptotic insults [23-25].

Taken together, the available *in vitro* and *in vivo* data highlight PKC α , PKC δ , and PKC ϵ as future candidates for targets in breast cancer therapy and as markers for disease prognosis. However, so far there is limited knowledge on the potential of the different isoforms as diagnostic and prognostic markers in breast cancer. This study sheds light on this issue by analyzing the expression levels of these PKC isoforms in primary breast

cancer tissue and our results indicate that $PKC\alpha$ is a potential marker of breast cancer aggressiveness.

Methods

Cell culture

All cell lines were obtained from ATCC. MCF-7, MDA-MB-231, and MDA-MB-468 breast cancer cells were maintained in RPMI 1640 medium (Sigma) supplemented with 10% fetal bovine serum (FBS; Invitrogen), 1 mM sodium pyruvate (PAA laboratories Gmbh), 100 IU/ml penicillin, and 100 µg/ml streptomycin (both Gibco). T47D cells were grown in DMEM supplemented with 10% FBS, 10 mM HEPES (PAA laboratories Gmbh), 100 IU/ml penicillin, and 100 µg/ml streptomycin. The media for MCF-7 and T47D cells were additionally supplemented 0.01 mg/ml insulin (Novo Nordisk A/S).

Transfections

For siRNA transfections, cells were seeded at 35-50% confluency and grown in complete medium without antibiotics for 24 hours. Cells were transfected for 48 hours using 4 μ /ml Lipofectamine 2000 (Invitrogen) and 40 nM siRNA (Invitrogen, table 1) in Optimem (Gibco) according to supplier's protocol.

Plasmid transfections were carried out for five hours replacing normal medium with Optimem containing 2 μ l/mL Lipofectamine 2000 and 2 μ g/mL DNA according to supplier's protocol. Plasmids encoding PKC constructs fused to enhanced green fluorescent protein (EGFP) have been described previously [26].

Tumor material

Cohort I originally consisted of tumors from 114 patients diagnosed with breast cancer at Umeå University Hospital and treated according to regional guidelines. The cohort is described in table 2. Due to lack of tumor material in the tissue microarray, 42-60 tumors could be analyzed, depending on the parameter investigated. Ki-67 had been classified in two groups <20% and >20% positive cells.

Cohort II included 512 consecutive breast cancer cases diagnosed at the department of Pathology, Malmö University Hospital, between 1988 and 1992. The cohort is described in table 2. Due to lack of tumor material in the tissue microarray, 223-263 tumors could be analyzed, depending on the parameter investigated. Ki-67 had been classified in three groups, 0-10%, 11-25% and 26-100% positive cells.

The cohorts represented a mix of all histological subtypes in proportions corresponding to the common incidence. The construction of tissue microarrays and clinicopathological properties of the cohorts have been described in detail elsewhere [27-32]. Ethical permissions were obtained from the Lund and Umeå Ethical Boards. The number of tumors that could not be evaluated for PKC expression due to lack of material is indicated as not evaluated in the tables.

Cell pellet arrays

Cells were washed in phosphate-buffered saline (PBS) and fixed for 25 minutes in 4% paraformaldehyde in PBS with Mayer's hematoxylin (5 μ l/ml) present during the last 5 minutes. The cells were pelleted, paraformaldehyde was removed and they were thereafter incubated over night in 70% ethanol followed by dehydration using increasing concentrations of ethanol and finally xylen. After dehydration, cell pellets were embedded in paraffin and arranged in a cell line array.

Immunohistochemistry

Sections (4 µm) of the paraffin blocks were dried, deparaffinized, rehydrated and microwave-treated in 1× target retrieval solution with high pH (DAKO). All sections were stained in a DAKO TechmateTM machine and visualized using DAB. The antibodies used were PKCa (1:2000), PKC δ (1:1000), PKC ϵ (1:400; all Santa Cruz Biotechnology, product numbers sc-208, sc-937 and sc-214), and Ki-67 (1:200; DAKO). All tissue microarray slides of a cohort were stained simultaneously with the same staining solutions ensuring identical conditions for each tumor. PKC stainings were scored according to cytoplasmic staining intensity were 0 representing lack of staining, 1 low staining, 2 moderate staining, and 3 strong staining. All cohorts were examined independently by two investigators and disconcordant results were re-evaluated.

For Ki-67 analyses of breast cancer cell lines, Ki-67 staining intensity was scored as negative-low or moderate-strong staining.

Sample preparation and Western blot

Cells were washed twice in ice-cold PBS and lyzed with RIPA buffer (10 mM Tris-HCl pH 7.2, 160 mM NaCl, 1% Triton-X 100, 1% sodium deoxycholate, 0.1% sodium dodecyl sulfate, 1 mM EDTA, 1 mM EGTA) supplemented with 40 μ l/ml complete protease inhibitor (Roche Applied Science) for 30 minutes on ice. Lysates were cleared by centrifugation at 14,000 × g for 10 minutes at 4°C.

Proteins were separated with SDS-PAGE and transferred to polyvinylidene difluoride membranes (Millipore). Membranes were blocked with PBS containing 0.05% tween and 5% non-fat milk, and probed with antibodies towards PKCδ (1:500), PKCε (1:500),

PKCα (1:3000), and actin (1:2000; MP Biomedicals, clone C4). Proteins were visualized with horseradish peroxidase-labeled secondary antibody (Amersham Biosciences) using the SuperSignal system (Pierce Chemical) as substrate. The chemiluminescence was detected with a CCD camera (Fuji Film).

Analysis of cell growth (WST-1 assay)

Cell were seeded at a density of 2000 cells per well in 96-well culture plates, and incubated for 24 hours. For cell line comparison, viable cell number was measured 24 and 48 hours after seeding. For experiments with inhibition or activation of PKC, 2 µM Gö6976 (Calbiochem) or equal volume DMSO, or 16 nM 12-*O*-tetradecanoylphorbol-13-acetate (TPA; Sigma) was added in complete medium (CM) or serum-free medium (SFM) 24 hours after seeding, and cells were incubated for 72 hours prior to estimation of viable cell number. The amount of viable cells was assessed by a WST-1 cell viability assay (Roche Applied Science). Absorbance was measured in an ELISA plate reader Antos 2020 (Antos Labtech Instruments).

Immunofluorescence and confocal microscopy

Immunofluorescence of PKCα was done as described [33] using Alexa Fluor 488conjugated secondary antibodies. Cells were examined with a Zeiss LSM 710 confocal system using standard settings for Alexa Fluor 488.

Cell cycle analysis

MDA-MB-231 cells were seeded at a density of 100,000 cells per 35 mm cell culture dish and transfected with siRNA. After transfection, cells were incubated in SFM or CM for 24 hours. Cells were trypsinized and fixed in 70% ethanol for 20 minutes at -20° C, washed in PBS, and incubated with a solution containing 3.5 μ M Tris-HCl pH 7.6, 10 mM NaCl, 50 μ g/ml propidium iodide (PI), 20 μ g/ml RNase, and 0.1% igepal CA-630 for 20 minutes on ice in order to label DNA. 10,000 events were acquired on the FL-2 channel for the PI signal. Sample acquisition and analyses were performed with CellQuest software (Becton Dickinson).

Wound healing assay

MDA-MB-231 or MCF-7 cells were seeded at a density of 150,000 cells per 35 mm cell culture dish and transfected with siRNA. After transfection, a scratch was made with a 200 µl pipette tip in a confluent area of the cell culture dish. Photographs of a selected area of each scratch were taken at indicated time points. For experiments with PKC inhibitors, MDA-MB-231 cells were seeded at a density of 350,000 cells per 35 mm cell culture dish and incubated for 24 hours before a scratch was made and PKC inhibitors were added. Photographs of a selected area of each scratching. The remaining wound area was measured using ImageJ software.

Statistics

For TMA analysis, correlations between variables were calculated using Pearson's twotailed significance test. Differences in distribution of various clinicopathological parameters in regard to PKC expression were also calculated using the χ^2 -test. The Kaplan-Meier analysis and the log rank test were used to illustrate differences between recurrence-free survival (RFS) and breast cancer-specific survival (BCSS) according to PKC expression. Cox regression proportional hazards models were used to estimate the impact of PKC α expression on breast cancer-specific survival in both univariate and multivariate analysis, adjusted for Nottingham histological grade (NHG), age, lymph node status, and tumor size of cohort II. For *in vitro* experiments, the significance of difference was assessed by analysis of variance (ANOVA) followed by Duncan's multiple range test. The difference was considered significant if the *p*-value was <0.05. All statistical calculations were performed using SPSS V.11.0.

Results

PKC expression in breast cancer tumors

Two cohorts of primary breast cancers (see experimental procedures for details) were analyzed for expression of PKC isoforms. Several batches of antibodies were tested to identify antibodies that did not cross-react with other isoforms. Cross reaction is a notorious problem for analyses of PKC isoforms. For PKC α , PKC δ , and PKC ϵ we had access to antibodies that did not cross-react with other isoforms (Figure 1A and B). As can be seen in Figure 1A there is only strong immunoreactivity in some of the cells where the cognate isoform has been overexpressed. In Figure 1B it can be seen that downregulation with siRNA abolishes or markedly diminishes the staining with the antibody towards the cognate isoform.

When present in a tumor, all of the PKC isoforms investigated were generally cytoplasmic and expressed in all of the tumor cells. Therefore, only the cytoplasmic staining intensity was accounted for in the analyses. Figure 1 C-J show examples of tumors with different staining intensities of PKC α , from negative to strong staining.

Initially a TMA of a smaller cohort (cohort I) was evaluated (Table 3 and 4). There was a significant correlation between PKC α staining intensity and lack of ER (p<0.001) and progesterone receptor (PR; p=0.002) as well as with tumor grade (p=0.001) and proliferation rate (Ki-67; p<0.001). PKC α levels did not correlate significantly with other clinicopathological parameters such as local or distal metastases (Table 3). For PKC δ and PKC ϵ there were no significant correlations with any of the clinicopathological parameters analyzed (Table 3).

A larger cohort (cohort II; Table 3 and 4) was then analyzed for expression levels of PKC α , since this isoform correlated to several clinicopatohological parameters in the first cohort, and PKCE, since our data somewhat contradict a published study [22]. The results from cohort II corroborated the findings in cohort I, where tumors with high PKCα levels had a higher proliferation rate (p<0.001), were ER (p<0.001) and PR-negative (p<0.001), and of a higher histological grade (p<0.001). Furthermore, a χ^2 analysis demonstrated that there is not an equal distribution of tumors with high and low PKCa-levels in the different histological subgroups in cohort II (Table 4). The skewed distribution is related to the tumors of medullary subtype. A tumor was defined as medullary if it fulfilled the following criteria: 1) lymphocytoplasmic reaction, 2) microscopic circumscription, 3) a syncytial growth pattern, and 4) poorly differentiated nuclear grade and high mitotic rate. For this group two out of twelve investigated tumors (17%) had high PKC α expression levels whereas the corresponding number for all evaluated breast cancers was five of 250 (2%). Only 8% of the medullary tumors were PKCa-negative, compared to 72% for all tumors. Thus, there is an overrepresentation of medullary carcinomas among tumors with high PKC α levels. For PKC ε , the findings were similar to cohort I, with no correlation to any relevant clinicopathological parameters.

PKCa expression is associated with poor prognosis

We next analyzed the relationship between the expression of PKC α and PKC ϵ and 10year survival of the patients (Figure 2). As illustrated in figure 2A, lower PKC α levels were associated with a significantly prolonged 10-year RFS (p=0.050). A similar trend was also seen for 10-year BCSS, but this did not reach statistical significance (Figure 2C; p=0.116). Since the majority of the tumors are negative for PKC α and the other groups (staining intensity 1-3) are small, a dichotomized variable defined as absent staining versus any staining was used for the same analyses of recurrence-free and breast cancer-specific survival. When dichotomized, PKC α positivity was associated with a non-significant trend towards a poorer 10-year RFS, figure 2B; p=0.074). However, patients with PKC α -negative tumors had a significantly improved 10-year BCSS compared to patients with PKC α -positive tumors (Figure 2D; p=0.016). We also performed a Cox regression proportional hazards analysis, demonstrating estimates of relative risks (RR) according to PKC α expression in univariate and multivariate analyses, adjusted for age at diagnosis, tumor size, NHG, node status and ER expression (Table 5). This revealed that the association between PKC α positivity and a poor 10-year BCSS was independent of established prognostic parameters (multivariate RR = 2.123, 95% CI 1.092 to 4.126, p=0.026).

Since a majority of the medullary carcinomas analyzed were PKC α -positive and patients with these tumors generally have good prognosis [34] we also examined 10-year BCSS after excluding patients with medullary carcinomas and found that PKC α remained an independent prognostic factor (Table 5).

Recurrence-free or breast cancer-specific 10-year survival was not significantly influenced by PKC ε expression, neither for all groups (staining intensity 0-3) nor when dichotomized into staining intensity 0-2 versus 3 (Figure 2E-H). A higher cutoff was chosen for dichotomization of PKC ε due to a smaller proportion of negative tumors.

PKC expression in breast cancer cell lines

To evaluate whether different breast cancer cell lines may represent the PKC isoform expression pattern in tumors, we measured PKC levels in four breast cancer cell lines (Figure 3A). High PKC α levels were seen in MDA-MB-231 cells compared to MCF-7 and MDA-MB-468 cells. PKC α was undetectable in T47D cells. PKC δ levels were higher in MCF-7 cells, whereas for PKC ϵ no major differences in expression levels could be noted.

To investigate whether the relationship between PKC α and a high proliferation rate observed in the tumors can be seen in cell lines as well, we examined the proliferation rate of the cell lines (Figure 3B and C). T47D, which had no detectable PKC α , was the least proliferative cell line, supporting the tumor data. On the other hand, for MDA-MB-231 which clearly had the highest PKC α levels, 97% of the cells had moderate or strong Ki-67 staining intensity (Figure 3B). For the cell lines with lower but detectable PKC α levels, MCF-7 and MDA-MB-468, the corresponding numbers were 88% and 91%, respectively. This did not differ significantly from MDA-MB-231 cells which would not be expected considering that the fraction of positive cells was close to 100% in these cell lines. The Ki-67 data were supported by the growth rate of the cell lines as estimated by a ratio of viable cell number after 48 h in culture with the number obtained after 24 h (Figure 3C).

PKCa is crucial for breast cancer cell proliferation under serum-free conditions.

To further examine whether PKCα is important for proliferation of breast cancer cells, the effects of a PKC activator and an inhibitor were studied (Figure 4A-C). Under serumfree conditions, activation of PKC by TPA was not sufficient to increase cell growth, as indicated by the amount of viable cells which represents the summative effect of proliferation and cell death. TPA did influence PKC α as seen by a relocation of PKC α to the plasma membrane and enrichment in the nucleus of both MCF-7 and MDA-MB-231 cells (Figure 4D). This indicates that PKC α is activated but the total levels of PKC α were also reduced following TPA treatment (Figure 4E) which complicates the estimation of the net effect of TPA on PKC α in the cells.

Gö6976, an inhibitor of classical PKCs, did not significantly reduce the number of cells indicating that PKC α activity is not critical for the growth of breast cancer cells under normal conditions.

PKC α has been shown to promote proliferation independently of its catalytic activity [35]. This fact together with the fact that both TPA and Gö6976 can have unspecific effects beside modulation of PKC α activity led us to more specifically analyze the role of PKC α by using siRNA oligonucleotides followed by analysis of cell cycle distribution. PKC α levels were downregulated in MDA-MB-231 cells with both siRNAs targeting PKC α and this did not influence the expression levels of the other PKC isoforms investigated (Figure 5A). Only one of the PKC α oligonucleotides (α #1) significantly influenced the cell cycle distribution when cells were grown in serum-containing complete medium (CM; Figure 5B). The oligonucleotide slightly decreased the amount of cells in S-phase compare to control. However, under serum-free medium (SFM) downregulation of PKC α with either siRNA led to significantly decreased amount of cells in S-phase compared to control cells (Figure 5C). The result indicates that PKC α plays an important role for proliferation of cells particularly under sub-optimal conditions. Under optimal growth conditions PKC α may be redundant for proliferation.

PKCa activity is necessary for migration of breast cancer cells

In breast cancer, PKCa has been associated with an increased ability of cells to migrate, since overexpression of PKC α has been shown to promote migration and metastasis of breast cancer cells [4, 36, 37]. Based on this we aimed at examining the importance of endogenous PKCa for breast cancer cell motility. Wound healing assays were performed with MDA-MB-231 cells in the absence or presence of PKC inhibitors (Figure 6A). Inhibition of classical PKCs (Gö6976) or classical and novel PKCs (GF109203X) suppressed the wound closure to 34% and 56%, respectively, of the DMSO treated cells, indicating that PKC activity is important for the ability of MDA-MB-231 cells to migrate. Since PKC inhibitors are isoform-unspecific, we downregulated PKCa with siRNA in MDA-MB-231 cells prior to a wound-healing assay, to more specifically study the role of PKC α in migration. However, downregulation of PKC α did not affect the migration of MDA-MB-231 cells (Figure 6B). MDA-MB-231 cells have high basal levels of PKCa compared to the other cell lines investigated in this study (Figure 3A). Transfection of these cells with siRNA targeting PKCa might not sufficiently reduce the PKCa levels. A comparison with MCF-7 cells (Figure 6C) actually demonstrates that following siRNA treatment of MDA-MB-231 cells the PKC α levels in these cells are roughly the same as in MCF-7 cells. Therefore, to obtain a more substantial PKC α depletion we downregulated PKC α in MCF-7 cells and thereafter performed a wound healing assay (Figure 6E). PKCa silencing in MCF-7 cells did not influence the other PKC isoforms investigated (Figure 6D). In these cells downregulation of PKC α significantly reduced migration into the wound to 55% of the control cells (Figure 6E). The results suggest that PKCα activity is crucial for migration of breast cancer cells *in vitro*.

Discussion

Several studies have suggested that DAG-sensitive PKC isoforms contribute to the progression of breast cancer and the malignant characteristics of breast cancer cells. In particular the PKC α , PKC δ , and PKC ϵ isoforms have been highlighted as potential targets for therapy of breast cancer or of specific subsets of the disease. This led us to design this study in which the expression of these PKC isoforms in primary breast cancer tumors has been examined to assess their utility as markers of tumor aggressiveness.

To substantiate the analysis, two different cohorts of primary breast cancer tumors were used, and the results from the cohorts were similar. We found significant correlations between PKC α levels and several markers of tumor aggressiveness including ER negativity. A correlation between PKC α levels and ER negativity has also been observed in a recent study [8] using 70 tumors from patients that had received systemic endocrine therapy. They also showed that high PKC α levels predicted a worse outcome in response to endocrine therapy, which is also supported in an earlier study with a smaller number of patients [7]. Our data, as demonstrated in two separate cohorts, firmly establish the relationship between PKC α and ER negativity. In addition, we identify a clear correlation between PKC α and PR negativity, and a positive correlation with tumor grade and high proliferation rate, further supporting the notion that PKC α expression is associated with parameters related to tumor aggressiveness. Finally, PKC α expression predicts a worse disease outcome with a significantly poorer 10-year breast cancer-specific survival for patients with primary tumors that were PKC α -positive. The results from the multivariate analysis further indicate that PKC α is an independent prognostic factor in breast cancer. However, other studies have indicated that PKC α levels actually are decreased in breast cancer compared to normal breast tissue [12, 13]. This may not necessarily contradict our findings since a vast majority of the cancer samples in our material were essentially PKC α -negative, which is in line with the notion that PKC α downregulation is a common event during breast cancer progression. Therefore, the published data, together with our results, suggest that most breast cancers are PKC α -negative, but that there are smaller subgroups with higher PKC α levels, displaying more aggressive clinicopathological features. If PKC α is to be used as a target for breast cancer therapy these data highlight the need to evaluate PKC α levels prior to such intervention.

In the largest cohort (II) there was also a significant association of PKC α levels with histological subtypes. Cancers with medullary histology were over-represented in the group with high PKC α levels. In addition, the only medullary cancer in cohort I was PKC α -positive.

In line with the tumor data demonstrating that PKC α levels correlate with features of aggressiveness, an association of PKC α with malignant features has also been seen in breast cancer cell lines. Increased levels of PKC α correlate to and can induce tamoxifen [38, 39] and multidrug [9, 10] resistance of ER-positive cell lines. In addition, overexpression of PKC α in MCF-7 cells leads to increased proliferation which is in line with our tumor data [4], but also make them more susceptible to apoptotic insults [40, 41]. We found that the cell line with no detectable PKC α expression, T47D, had a lower percentage of Ki-67-positive cells and slower growth rate than the other cell lines, similar to the tumor data. However, the cell line with the highest PKC α expression, MDA-MB-231, only had marginally higher Ki-67 positivity compared with cell lines with detectable

but much lower expression. This may be related to the fact that the fraction of Ki-67positive cells was high (97%) and could not be further elevated.

Neither inhibition of PKC α in the presence of serum nor PKC activation under serumstarvation influenced the growth of the cell lines in a manner that would support an essential role for PKC α for breast cancer cell growth. A recent paper has shown that the PKC α protein, but not activity, is essential for glioma cell proliferation [35]. We found that downregulation of PKC α with siRNA caused a modest effect on cell cycle distribution of MDA-MB-231 cells grown in complete medium. However, under serumfree conditions, PKC α silencing clearly reduced the amount of proliferating cells suggesting that, as in glioma cells, the PKC α protein, but maybe not its catalytic activity, supports proliferation under sub-optimal conditions.

Studies *in vitro* have demonstrated that increasing the PKC α expression in MCF-7 cells makes them more migratory in response to PKC activation [42]. Our experiments with cell lines indicate that PKC activity supports migration of MDA-MB-231 cells. However, downregulation of PKC α with siRNA in this cell line did not affect cell migration. MDA-MB-231 cells have high basal expression levels of PKC α and incomplete downregulation of PKC α may explain the lack of effect on cell migration. This assumption is supported by the more efficiently suppressed wound healing upon downregulation of PKC α in MCF-7 cells. Our data therefore indicate that PKC α activity is important for migration of breast cancer cells, in line with previous findings using PKC α overexpression. A migratory propensity might facilitate the metastasation of a tumor. However, our tumor data did not support a role for PKC α in dissemination. There was no correlation between PKC α expression and nodal or distant metastases. Thus, the effects of PKC α on migration may primarily be of importance *in vitro*.

For PKC δ there is less information regarding expression levels in primary tumors. One study has shown that low levels of PKC δ , particularly in combination with high PKC α , predict resistance to endocrine therapy [8]. In this study, we could not observe any significant associations between PKC δ expression and relevant clinicopathological parameters. Thus, altered PKC δ expression does not seem to be a prerequisite for breast cancer progression.

PKC ε has been proposed to be a marker of aggressive breast cancer since its expression was reported to be elevated in hormone receptor-negative breast cancers with high tumor grade and *HER2* amplification [22]. However, a relationship between PKC ε expression and tumor grade could not be confirmed in this study, suggesting that it might not be apparent in cohorts representing all histological subtypes of breast cancer. It is possible that the level of PKC ε is a marker of aggressiveness in more defined subgroups of breast cancer.

Conclusions

In conclusion, our findings demonstrate that expression of PKC α correlates to both ER and PR negativity as well as high histological grade and proliferation rate in breast cancer. An important role for PKC α in breast cancer cell proliferation was also observed *in vitro*. PKC α expression is not associated with metastasis in breast cancer samples, but PKC α activity supports migration of breast cancer cells *in vitro*. Importantly, PKC α expression predicts for a poorer 10-year breast cancer-specific survival, independently of established prognostic parameters. Thus, we propose that PKC α might be a useful marker for prognostication and treatment stratification in breast cancer.

Competing interests

The authors declare that they have no competing interests.

Author's contributions

GKL evaluated clinical samples, performed statistical analyses, participated in the design of and did most of the experimental work, and assembled the drafts of the manuscript. LC did some experimental work.

IOZ took part in the evaluation of the antibodies and clinical samples.

GL supervised the analysis of cohort I and participated in interpretative discussions.

KJ supervised the analysis of cohort II and statistical analyses of patient data, and helped draft the manuscript.

CL conceived of the study, participated in the design of the experimental work and helped draft the manuscript.

All authors read and approved the final manuscript.

Acknowledgements

We are grateful for excellent technical assistance from Elise Nilsson. This work was supported by grants from The Swedish Cancer Society, The Swedish Research Council, The Children's Cancer Foundation of Sweden, Malmö University Hospital Research Funds, and the Kock, Crafoord, Ollie and Elof Ericsson and Gunnar Nilsson Foundations.

References

- 1. Vargo-Gogola T, Rosen JM: Modelling breast cancer: one size does not fit all. *Nat Rev Cancer* 2007, **7:**659-672.
- 2. Sorlie T, Perou CM, Tibshirani R, Aas T, Geisler S, Johnsen H, Hastie T, Eisen MB, van de Rijn M, Jeffrey SS, et al: Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. *Proc Natl Acad Sci U S A* 2001, **98**:10869-10874.
- 3. Newton AC: Protein kinase C: structural and spatial regulation by phosphorylation, cofactors, and macromolecular interactions. *Chem Rev* 2001, **101**:2353-2364.
- Ways DK, Kukoly CA, deVente J, Hooker JL, Bryant WO, Posekany KJ, Fletcher DJ, Cook PP, Parker PJ: MCF-7 breast cancer cells transfected with protein kinase C-α exhibit altered expression of other protein kinase C isoforms and display a more aggressive neoplastic phenotype. J Clin Invest 1995, 95:1906-1915.
- Chisamore MJ, Ahmed Y, Bentrem DJ, Jordan VC, Tonetti DA: Novel antitumor effect of estradiol in athymic mice injected with a T47D breast cancer cell line overexpressing protein kinase Cα. Clin Cancer Res 2001, 7:3156-3165.
- 6. Tonetti DA, Chisamore MJ, Grdina W, Schurz H, Jordan VC: Stable transfection of protein kinase C α cDNA in hormone-dependent breast cancer cell lines. *Br J Cancer* 2000, 83:782-791.
- Tonetti DA, Morrow M, Kidwai N, Gupta A, Badve S: Elevated protein kinase C α expression may be predictive of tamoxifen treatment failure. *Br J Cancer* 2003, 88:1400-1402.
- 8. Assender JW, Gee JM, Lewis I, Ellis IO, Robertson JF, Nicholson RI: Protein kinase C isoform expression as a predictor of disease outcome on endocrine therapy in breast cancer. *J Clin Pathol* 2007, **60**:1216-1221.
- 9. Gill PK, Gescher A, Gant TW: Regulation of MDR1 promoter activity in human breast carcinoma cells by protein kinase C isozymes alpha and theta. *Eur J Biochem* 2001, **268**:4151-4157.
- Blobe GC, Sachs CW, Khan WA, Fabbro D, Stabel S, Wetsel WC, Obeid LM, Fine RL, Hannun YA: Selective regulation of expression of protein kinase C (PKC) isoenzymes in multidrug-resistant MCF-7 cells. Functional significance of enhanced expression of PKC α. J Biol Chem 1993, 268:658-664.
- Roychowdhury D, Lahn M: Antisense therapy directed to protein kinase C-α (Affinitak, LY900003/ISIS 3521): potential role in breast cancer. Semin Oncol 2003, 30:30-33.
- 12. Ainsworth PD, Winstanley JH, Pearson JM, Bishop HM, Garrod DR: Protein kinase C α expression in normal breast, ductal carcinoma in situ and invasive ductal carcinoma. *Eur J Cancer* 2004, **40**:2269-2273.
- Kerfoot C, Huang W, Rotenberg SA: Immunohistochemical analysis of advanced human breast carcinomas reveals downregulation of protein kinase C α. J Histochem Cytochem 2004, 52:419-422.

- 14. Zeidan YH, Wu BX, Jenkins RW, Obeid LM, Hannun YA: A novel role for protein kinase Cδ-mediated phosphorylation of acid sphingomyelinase in UV light-induced mitochondrial injury. *Faseb J* 2008, 22:183-193.
- McCracken MA, Miraglia LJ, McKay RA, Strobl JS: Protein kinase C δ is a prosurvival factor in human breast tumor cell lines. *Mol Cancer Ther* 2003, 2:273-281.
- Nabha SM, Glaros S, Hong M, Lykkesfeldt AE, Schiff R, Osborne K, Reddy KB: Upregulation of PKC-δ contributes to antiestrogen resistance in mammary tumor cells. Oncogene 2005, 24:3166-3176.
- Alonso-Escolano D, Medina C, Cieslik K, Radomski A, Jurasz P, Santos-Martinez MJ, Jiffar T, Ruvolo P, Radomski MW: Protein kinase C δ mediates platelet-induced breast cancer cell invasion. J Pharmacol Exp Ther 2006, 318:373-380.
- Kiley SC, Clark KJ, Duddy SK, Welch DR, Jaken S: Increased protein kinase C δ in mammary tumor cells: relationship to transformation and metastatic progression. Oncogene 1999, 18:6748-6757.
- Kiley SC, Clark KJ, Goodnough M, Welch DR, Jaken S: Protein kinase C δ involvement in mammary tumor cell metastasis. *Cancer Res* 1999, 59:3230-3238.
- Grossoni VC, Falbo KB, Kazanietz MG, de Kier Joffe ED, Urtreger AJ: Protein kinase C δ enhances proliferation and survival of murine mammary cells. *Mol Carcinog* 2007, 46:381-390.
- 21. Lønne GK, Masoumi KC, Lennartsson J, Larsson C: Protein kinase Cδ supports survival of MDA-MB-231 breast cancer cells by suppressing the ERK1/2 pathway. *J Biol Chem* 2009, **284**:33456-33465.
- Pan Q, Bao LW, Kleer CG, Sabel MS, Griffith KA, Teknos TN, Merajver SD: Protein kinase C ε is a predictive biomarker of aggressive breast cancer and a validated target for RNA interference anticancer therapy. *Cancer Res* 2005, 65:8366-8371.
- 23. Lu D, Sivaprasad U, Huang J, Shankar E, Morrow S, Basu A: **Protein kinase C-ε** protects MCF-7 cells from TNF-mediated cell death by inhibiting Bax translocation. *Apoptosis* 2007.
- 24. Sivaprasad U, Shankar E, Basu A: **Downregulation of Bid is associated with PKCe-mediated TRAIL resistance.** *Cell Death Differ* 2007, **14**:851-860.
- 25. Lu D, Huang J, Basu A: Deregulation of PKB influences antiapoptotic signaling by PKC in breast cancer cells. *Int J Oncol* 2004, **25:**671-676.
- 26. Zeidman R, Lofgren B, Pahlman S, Larsson C: **PKCe**, via its regulatory domain and independently of its catalytic domain, induces neurite-like processes in neuroblastoma cells. *J Cell Biol* 1999, **145**:713-726.
- 27. Nielsen NH, Arnerlov C, Emdin SO, Landberg G: Cyclin E overexpression, a negative prognostic factor in breast cancer with strong correlation to oestrogen receptor status. *Br J Cancer* 1996, **74**:874-880.
- 28. Nielsen NH, Emdin SO, Cajander J, Landberg G: **Deregulation of cyclin E and D1 in breast cancer is associated with inactivation of the retinoblastoma protein.** *Oncogene* 1997, **14:**295-304.

- 29. Roos G, Nilsson P, Cajander S, Nielsen NH, Arnerlov C, Landberg G: Telomerase activity in relation to p53 status and clinico-pathological parameters in breast cancer. *Int J Cancer* 1998, **79:**343-348.
- 30. Svensson S, Jirstrom K, Ryden L, Roos G, Emdin S, Ostrowski MC, Landberg G: ERK phosphorylation is linked to VEGFR2 expression and Ets-2 phosphorylation in breast cancer and is associated with tamoxifen treatment resistance and small tumours with good prognosis. Oncogene 2005, 24:4370-4379.
- 31. Borgquist S, Holm C, Stendahl M, Anagnostaki L, Landberg G, Jirstrom K: Oestrogen receptors α and β show different associations to clinicopathological parameters and their co-expression might predict a better response to endocrine treatment in breast cancer. J Clin Pathol 2008, 61:197-203.
- 32. Helczynska K, Larsson A-M, Holmquist Mengelbier L, Bridges E, Fredlund E, Borgquist S, Landberg G, Pahlman S, Jirstrom K: **Hypoxia-inducible factor-2α correlates to distant recurrence and poor outcome in invasive breast cancer**. *Cancer Res* 2008, **68**:9212-9220.
- 33. Raghunath A, Ling M, Larsson C: The catalytic domain limits the translocation of protein kinase C α in response to increases in Ca2+ and diacylglycerol. *Biochem J* 2003, **370**:901-912.
- 34. Malyuchik SS, Kiyamova RG: Medullary breast carcinoma. *Exp Oncol* 2008, **30**:96-101.
- 35. Cameron AJ, Procyk KJ, Leitges M, Parker PJ: **PKC** α protein but not kinase activity is critical for glioma cell proliferation and survival. *Int J Cancer* 2008, **123:**769-779.
- 36. Ng T, Shima D, Squire A, Bastiaens PI, Gschmeissner S, Humphries MJ, Parker PJ: **PKC** α regulates β 1 integrin-dependent cell motility through association and control of integrin traffic. *Embo J* 1999, **18**:3909-3923.
- 37. Abeyweera TP, Chen X, Rotenberg SA: **Phosphorylation of α 6-tubulin by protein kinase Cα activates motility of human breast cells.** *J Biol Chem* 2009.
- 38. Frankel LB, Lykkesfeldt AE, Hansen JB, Stenvang J: Protein Kinase C α is a marker for antiestrogen resistance and is involved in the growth of tamoxifen resistant human breast cancer cells. Breast Cancer Res Treat 2007, 104:165-179.
- 39. Lin X, Yu Y, Zhao H, Zhang Y, Manela J, Tonetti DA: Overexpression of PKCα is required to impart estradiol inhibition and tamoxifen-resistance in a T47D human breast cancer tumor model. Carcinogenesis 2006, 27:1538-1546.
- 40. Yde CW, Gyrd-Hansen M, Lykkesfeldt AE, Issinger OG, Stenvang J: Breast cancer cells with acquired antiestrogen resistance are sensitized to cisplatininduced cell death. *Mol Cancer Ther* 2007, **6**:1869-1876.
- 41. de Vente JE, Kukoly CA, Bryant WO, Posekany KJ, Chen J, Fletcher DJ, Parker PJ, Pettit GJ, Lozano G, Cook PP, et al.: Phorbol esters induce death in MCF-7 breast cancer cells with altered expression of protein kinase C isoforms. Role for p53-independent induction of gadd-45 in initiating death. J Clin Invest 1995, 96:1874-1886.

42. Parsons M, Keppler MD, Kline A, Messent A, Humphries MJ, Gilchrist R, Hart IR, Quittau-Prevostel C, Hughes WE, Parker PJ, Ng T: Site-directed perturbation of protein kinase C- integrin interaction blocks carcinoma cell chemotaxis. *Mol Cell Biol* 2002, 22:5897-5911.

Legends to figures

Figure 1

Immunohistochemical stainings of primary breast cancer and validation of antibody specificity. MCF-7 cells were transfected with vectors encoding PKC α (α), PKC δ (δ), or PKC ϵ (ϵ) (A) and MDA-MB-231 cells were mock-treated or transfected with siRNA targeting PKC α (α), PKC δ (δ), or PKC ϵ (ϵ) (B). Pellets of transfected cells were arranged in a cell line array and immunohistochemistry was performed with antibodies towards indicated PKC isoforms. (C-J) Examples of immunohistochemical staining of PKC α in breast cancer specimens from cohort II showing negative (C and G), low (D and H), moderate (E and I), and intense (F and J) staining with 20× magnification (C-F) and with 40× magnification (G-J).

Figure 2

Recurrence-free survival and breast cancer-specific survival according to PKCα and PKCε expression. Kaplan-Meier estimates of 10 year recurrence-free survival (A-B and E-F) and breast cancer-specific survival (C-D and G-H) according to PKCα (A-D) and PKCε (E-H) expression.

Figure 3

PKC expression and proliferation of breast cancer cell lines. Western blots show expression levels of PKCα, PKCδ, and PKCε in T47D, MCF-7, MDA-MB-231, and MDA-MB-468 breast cancer cells (A). Cell pellets of proliferating cells were immunohistochemically stained with anti-Ki-67. At least 200 cells per cell line were

scored for negative-weak or strong staining intensity (B). The viable cell number was measured by WST-1 assay after 24 and 48 hours in culture. Ratio 48 h:24 h was used as an indicator of cell proliferation (C). For all experiments, cells were grown in complete medium. Western blots are representative of three independent experiments. Data in B and C are mean \pm SEM, n = 3. Asterisks indicate statistically significant differences (* p<0.05 and ** p<0.01) compared to other cell lines (B and C).

Figure 4

Cell growth analysis after activation or inhibition of PKC. T47D (A), MCF-7 (B), and MDA-MB-231 (C) breast cancer cells were incubated in the absence or presence of 16 nM TPA in serum free medium (SFM) or 2 μ M Gö6976, GF109203X, or equal volume of DMSO in complete growth medium (CM) for 72 hours prior to WST-1 assay. Data (mean \pm SEM, n = 3) represent the amount of viable cells expressed as percent of viable cells obtained under control conditions. The localization of PKC α in control or TPA-treated MCF-7 and MDA-MB-231 cells was examined with immunofluorescence followed by confocal microscopy (D). The levels of PKC α following treatment with TPA or Gö6976 were analyzed by Western blot (E).

Figure 5

Cell cycle distribution of MDA-MB-231 cells with downregulated PKCa. MDA-MB-231 cells were transfected with two different siRNAs targeting PKCa (α #1 and α #2) or a control oligonucleotide. After transfection, cells were incubated with complete medium (CM) or serum-free medium (SFM) for 24 hours. Adherent cells were thereafter

subjected to Western blot (A) or propidium iodide staining and flow cytometry (B and C). Western blots are representative of three independent experiments. Data in B and C (mean \pm SEM, n = 3) show the percentage of cells in s-phase.

Figure 6

Migration of MDA-MB-231 and MCF-7 cells. MDA-MB-231 cells were subjected to wound-healing assay in presence or absence of PKC inhibitors (A) or after transfection with siRNA targeting PKC α (α) or a control oligonucleotide (c) (B). PKC α was downregulated in MCF-7 cells (C and D) before wound-healing assay was performed (E). Data (mean \pm SEM, n = 4) show wound closure compared to closure in control conditions. Asterisks indicate statistically significant differences (* *p*<0.05 and ** *p*<0.01) compared to control conditions.

Tables

Table 1 siRNA oligonucleotides

siRNA	Oligonucleotides
Control	GACAGUUGAACGUCGAUUUGCAUUG
PKCa #1	CCGAGUGAAACUCACGGACUUCAAU
PKCa #2	CCAUCGGAUUGUUCUUCUUCAUAA
ΡΚϹδ	CCAAGGUGUUGAUGUCUGUUCAGUA
ΡΚϹε	CACAAGUUCGGUAUCCACAACUACA

Table 2 Characteristics of the cohorts investigated

Cohort	Ι	II
Number of patients	114	512
Age at diagnosis, median (range)	60 (30-80)	65 (27-96)
Tumor size (mm), median (range)	22 (8-100)	16 (1-100)
Nodal status		
Positive	53	168
Negative	48	298
Missing	13	58
ER-status		
Positive	82	417
Negative	31	72
Missing	1	35

Table 3
Associations between PKCa, PKCb, and PKCc intensity and clinicopathological
variables in cohort I and II.

Cohort	Ι			П	
Variable	PKCa	РКСб	ΡΚϹε	РКСа	ΡΚϹε
NHG					
ρ	0.401	-0.049	-0.029	0.247	-0.011
p	0.001	0.719	0.833	<0.001	0.863
n	60	57	55	249	263
Estrogen receptor					
ρ	-0.482	0.210	-0.016	-0.334	0.068
p	<0.001	0.120	0.907	<0.001	0.273
n	59	56	54	244	261
Progesterone receptor					
ρ	-0.402	-0.081	-0.216	-0.280	0.062
p	0.002	0.554	0.120	<0.001	0.342
n	58	55	53	223	240
Ki-67 positivity					
ρ	0.535	0.229	0.280	0.295	-0.056
p	<0.001	0.144	0.069	<0.001	0.363
n	46	42	43	244	261
Nodal status					
ρ	-0.027	0.095	-0.002	0.029	0.028
р	0.842	0.495	0.987	0.666	0.667
n	56	54	52	224	233
Distant metastasis					
ρ	0.159	0.162	-0.064	0.090	0.037
р	0.226	0.227	0.642	0.159	0.556
n	60	57	55	247	261

 ρ : Pearson's correlation coefficient. p < 0.05 in bold. n: number of tumor samples.

Table 4 Associations between PKCα, PKCδ, and PKCε intensity and histological type in cohort I and II.

Cohort I		Ductal	Lobular	Medullary	Mucinous		
PKCa intensity	0	25	3	0	2		
p*=0.295	1	20	0	1	0		
Ι	2	4	0	0	0		
	3	4	0	1	0		
	n.e.	42	3	0	2		
						-	
PKC ₀ intensity	0	12	0	0	1		
p*=0.850	1	33	1	2	1		
	2	7	0	0	0		
	3	0	0	0	0		
	n.e.	43	5	0	2		
DKC a intensity	0	0	0	0	0	-	
$r K C \epsilon$ intensity $r^* = 0.800$	1	12	0	1	0		
p*=0.800	1	20	1	1	0		
	23	0	1	1	2		
	ne	15	5	0	2		
	п.с.	45	5	0	2	-	
Cohort II		Ductal	Lobular	Medullary	Mucinous	Tubular	Mixed
PKCa intensity	0	124	17	1	10	14	13
p*< 0.001	1	34	4	5	2	3	3
P 100001	2	10	0	4	0	0	1
	3	3	0	2	0	0	0
	n.e.	168	52	3	5	17	17
	0	2	1			0	0
PKCe intensity	0	2	1	0	0	0	0
p*=0.328	1	58	9	/	4	5	5
	2	90	19	4	/	9	/
	3	41	5	1	1	1	1
		1/0	20	2	~	10	1 7

 $^{*}\chi^{2}$ -test, p<0.05 in bold, n.e.=not evaluated

Table 5

Cox univariate and multivariate analysis of breast cancer-specific survival in cohort II according to PKC α expression in all patients and when patients with medullary cancer are excluded.

	All patients		Medullary cancers excluded		
	RR (95%CI)	p value	RR (95%CI)	p value	
	Univariate		Univariate		
PKCα negative	1.00		1.00		
PKCa positive	2.105 (1.148 to 3.862)	0.016	1.995 (1.069 to 3.721)	0.030	
PKCa negative	<i>Multivariate</i> 1.00		<i>Multivariate</i>		
PKCa positive	2.123 (1.092 to 4.126)	0.026	1.978 (0.984 to 3.975)	0.055	

Multivariate analysis adjusted for Nottingham histological grade (I-II/III), age (continuous), nodal status (0/1), tumor size (continuous), and ER (< or > 10%).



















MCF-7

Π



MDA-MB-231







Figure 4











С

Figure 5

α#1

α #2









Figure 6