

This Provisional PDF corresponds to the article as it appeared upon acceptance. Copyedited and fully formatted PDF and full text (HTML) versions will be made available soon.

## Expression levels of uridine 5'-diphospho-glucuronosyltransferase genes in breast tissue from healthy women are associated with mammographic density

*Breast Cancer Research* 2010, **12**:R65 doi:10.1186/bcr2632

Vilde D Haakensen (vilde.drageset.haakensen@rr-research.no)  
Margarethe Biong (Margarethe.Biong@rr-research.no)  
Ole Christian Lingjaerde (ole@ifi.uio.no)  
Marit Muri Holmen (Marit.Muri.Holmen@radiumhospitalet.no)  
Jan Ole Frantzen (jan.ole.frantzen@unn.no)  
Ying Chen (ying.chen@siv.no)  
Dina Navjord (Dina.Navjord@sykehuset-innlandet.no)  
Linda Romundstad (linda.romundstad@sb-hf.no)  
Torben Luders (torben.luders@medisin.uio.no)  
Ida K Bukholm (i.r.k.bukholm@medisin.uio.no)  
Hiroko K Solvang (Hiroko.Solvang@rr-research.no)  
Vessela N Kristensen (vessela@ulrik.uio.no)  
Giske Ursin (giske.urisn@medisin.uio.no)  
Anne-Lise Borresen-Dale (a.l.borresen-dale@medisin.uio.no)  
Aslaug Helland (Aslaug.Helland@rr-research.no)

**ISSN** 1465-5411

**Article type** Research article

**Submission date** 22 February 2010

**Acceptance date** 27 August 2010

**Publication date** 27 August 2010

**Article URL** <http://breast-cancer-research.com/content/12/4/R65>

This peer-reviewed article was published immediately upon acceptance. It can be downloaded, printed and distributed freely for any purposes (see copyright notice below).

Articles in *Breast Cancer Research* are listed in PubMed and archived at PubMed Central.

For information about publishing your research in *Breast Cancer Research* go to

© 2010 Haakensen *et al.*, licensee BioMed Central Ltd.

This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



# Expression levels of uridine 5'-diphosphoglucuronosyltransferase genes in breast tissue from healthy women are associated with mammographic density

Vilde D Haakensen<sup>1,2</sup>, Margarethe Biong<sup>1</sup>, Ole Christian Lingjærde<sup>3,4</sup>, Marit Muri Holmen<sup>5</sup>, Jan Ole Frantzen<sup>6</sup>, Ying Chen<sup>7</sup>, Dina Navjord<sup>8</sup>, Linda Romundstad<sup>9</sup>, Torben Lüders<sup>10</sup>, Ida K Bukholm<sup>2, 11</sup>, Hiroko K Solvang<sup>1</sup>, Vessela N Kristensen<sup>1, 2, 10</sup>, Giske Ursin<sup>12, 13</sup>, Anne-Lise Børresen-Dale<sup>1, 2</sup>, Åslaug Helland<sup>1, 2, 14 \*</sup>

<sup>1</sup> Department of Genetics, Institute for Cancer Research, Oslo University Hospital Radiumhospitalet, Oslo, Montebello, NO-0310, Norway

<sup>2</sup> Institute for Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, NO-0315, Norway

<sup>3</sup> Biomedical Research Group, Department of Informatics, University of Oslo, Oslo, NO-0315, Norway

<sup>4</sup> Centre for Cancer Biomedicine, University of Oslo, Oslo, NO-0315, Norway

<sup>5</sup> Department of Radiology, Oslo University Hospital Radiumhospitalet, Oslo, NO-0310, Norway

<sup>6</sup> Department of Radiology, University Hospital of North Norway, Tromsø, NO- 9038, Norway

<sup>7</sup> Department. of Pathology, Vestfold Hospital, Halfdan Wilhelmsens Alle' 17, Tønsberg, NO-3103, Norway

<sup>8</sup> Department of Radiology, Innlandet Hospital, Brummundal, NO-2381, Norway

<sup>9</sup> Department of Radiology, Buskerud Hospital, Drammen, NO-3004, Norway

<sup>10</sup> Department for Clinical Molecular Biology (EpiGen), Institute for Clinical Medicine, Akershus University Hospital, University of Oslo, Oslo, NO-0315, Norway

<sup>11</sup> Department of Surgery, Akerhus University Hospital, Lørenskog, 1478, Norway

<sup>12</sup> Department of Nutrition, School of Medicine, University of Oslo, Oslo, NO-0315, Norway

<sup>13</sup> Department of Preventive Medicine, University of Southern California Keck School of Medicine, Los Angeles, CA 90033, USA

<sup>14</sup> Department of Oncology, Oslo University Hospital Radiumhospitalet, Oslo, NO-0310, Norway

\*Corresponding author: [Aslaug.helland@rr-research.no](mailto:Aslaug.helland@rr-research.no)

# **Abstract**

## **Introduction**

Mammographic density (MD), as assessed from film screen mammograms, is determined by the relative content of adipose, connective and epithelial tissue in the female breast. In epidemiological studies, a high percentage of MD confers a four to six fold risk elevation of developing breast cancer, even after adjustment for other known breast cancer risk factors. However, the biologic correlates of density are little known.

## **Methods**

Gene expression analysis using whole genome arrays was performed on breast biopsies from 143 women; 79 women with no malignancy (healthy women) and 64 newly diagnosed breast cancer patients, both included from mammographic centres. Percent MD was determined using a previously validated, computerized method on scanned mammograms. Significance analysis of microarrays (SAM) was performed to identify genes influencing MD and a linear regression model was used to assess the independent contribution from different variables to MD.

## **Results**

SAM-analysis identified 24 genes differentially expressed between samples from breasts with high and low MD. These genes included three uridine 5'-diphospho-glucuronosyltransferase (*UGT*) genes and the oestrogen receptor gene (*ESR1*). These

genes were down-regulated in samples with high MD compared to those with low MD.

The UGT gene products, which are known to inactivate oestrogen metabolites, were also down-regulated in tumour samples compared to samples from healthy individuals.

Several single nucleotide polymorphisms (SNPs) in the *UGT* genes associated with the expression of *UGT* and other genes in their vicinity were identified.

## **Conclusions**

Three UGT enzymes were lower expressed both in breast tissue biopsies from healthy women with high MD and in biopsies from newly diagnosed breast cancers. The

association was strongest amongst young women and women using hormonal therapy.

UGT2B10 predicts MD independently of age, hormone therapy and parity. Our results indicate that down-regulation of *UGT* genes in women exposed to female sex hormones is associated with high MD and might increase the risk of breast cancer.

## Introduction

Breast cancer is a common disease in women. Knowledge about the first steps in tumour initiation is important for early detection. However, the exact mechanisms of tumour initiation are still unknown.

Mammographic density (MD), captured on film screen mammograms, refers to the content and architectural structure of the adipose, connective and epithelial tissues in the female breast [1]. In epidemiological studies, a high percentage of MD confers a four to six fold elevated risk of developing breast cancer [1-3] and has been proposed as a possible surrogate marker for the disease[4]. The relative risk associated with mammographic densities remains at this magnitude even after adjustment for all other known breast cancer risk factors. Breasts with high MD have greater tissue cellularity and more tissue collagen [5]. Still, little is known as to how MD confers the increased breast cancer risk. MD is to a large degree an inherited trait, although it is also influenced by environmental factors, hormone therapy being an evident example [6]. The genetic factors determining the inheritability are largely unknown.

In order to elucidate how MD increases risk of breast cancer; we searched for the biological correlates to MD. Gene expression analysis on biopsies from breasts of healthy women with various degrees of MD was performed. The gene expression profiles represent the gene activity of the different cell types in the biopsy, producing a fingerprint of the breast tissue within the biopsy of that particular woman.

The breast is an oestrogen-sensitive organ. MD varies with levels of female hormones, and is reduced after menopause. The uridine 5'-diphospho-glucuronosyltransferase (*UGT*) genes encode enzymes inactivating several endogenous and exogenous compounds, including sex hormones (Figure 1) [7]. *UGT1A1* is known to be responsible for the glucuronidation of bilirubin, but is also shown to glucuronidate catechol oestrogens [8,9]. Polymorphisms in this gene have previously been linked to MD in premenopausal women [10]. *UGT2B7* is known to conjugate oestrone, one of the active oestradiol metabolites. This enzyme has previously been found down-regulated in tumour tissue compared with non-malignant tissue leading to the conclusion that *UGT* expression could lead to promotion of carcinogenesis [11] but there are no reports on this gene in relation to MD in the literature. Less is known about the other *UGT2B* genes, although here is extensive structural homology. We will use the *UGT* genes as a term describing three *UGT2B* genes significantly down-regulated in our analyses (*UGT2B7*, 10 and 11). Other *UGT* genes are specified in the text. In this study we analysed biopsies from breasts of healthy women and found genes whose expression is associated with MD.

## Material and methods

### Subjects

The women included in this study had all attended one of 6 breast diagnostic centres in Norway which are part of the governmentally funded National Breast Cancer Screening Program between 2002-2007 [12]. Women were eligible if they did not currently use anticoagulants, did not have breast implants and were not currently pregnant or lactating. A total of 186 women were recruited to the study; 120 healthy women with no malignant disease but some visible density in the mammograms, in the following referred to as healthy women, and 66 women with a newly diagnosed breast cancer. Of these, quality tested expression data were obtained from biopsies from 79 healthy women and 64 breast cancer patients.

The women were either referred to a breast diagnostic centre for a second look due to some irregularity of the initial screening mammogram (n=69), or due to clinical findings (n=83). For 34 women the type of referral was unknown.

The women provided information about height, weight, parity, hormone therapy use and family history of breast cancer. Two breast biopsies and three blood samples were collected from each woman. All women provided signed informed consent. The study was approved by the local ethical committee and local authorities (IRB approval no S-02036).

## **Core biopsies**

Two breast biopsies were obtained from each woman with a 14 gauge needle, for RNA- and DNA-extraction. In healthy women, the biopsies were taken from an area with no visible pathology, but with some MD to ensure that the biopsies did not contain only fatty tissue, which yields little RNA. The sampling was guided by ultrasound. At one hospital, six of the biopsies from breasts of healthy women were collected from a benign lesion (mostly fibroadenomas). For the cancer patients, all biopsies were taken from the tumour. The tissue was either fresh-frozen at  $-80^{\circ}\text{C}$  or soaked in ethanol and RNAlater (Ambion, Austin, TX), transported and subsequently stored at  $-20^{\circ}\text{C}$ .

## **Pathology**

The haematoxylin eosinofil sections from the tumours of the breast cancer patients were evaluated locally and then re-evaluated by one pathologist (YC). Information about tumour size, histological grade and type, oestrogen and progesterone receptor status, HER2 status and sentinel node status was recorded and entered into a database managed by the Office for Clinical Research at Oslo University Hospital, Radiumhospitalet. Pathology evaluations were not available for the biopsies from breasts of healthy women.

## **RNA-expression analysis**

Homogenization, cell lysis and RNA extraction was performed using the RNeasy Mini Protocol (Qiagen, Valencia, CA). RNA quality was controlled by Agilent 2100 Bioanalyzer (Agilent Technologies, Palo Alto, CA) and concentration was determined using NanoDrop ND-1000 spectrophotometer (Thermo Scientific, Wilmington DE). A total of 40 samples, mostly from normal breast tissue, were excluded from further analyses due to low RNA amount (<10ng) or poor RNA quality. RNA was then amplified and labelled using the Agilent Low RNA input Fluorescent Linear Amplification Kit Protocol. Amplified tumour RNA was labelled by Cy5 (Amersham Biosciences, Little Chalfont, England) and amplified RNA from Universal Human total RNA (Stratagene, La Jolla, CA) was labelled by Cy3 (Amersham Biosciences, Little Chalfont, England). RNA from the remaining 146 biopsies was further hybridized on Agilent Human Whole Genome Oligo Microarrays (G4110A) (Agilent Technologies, Santa Clara, CA). Three arrays had to be excluded due to poor quality leaving data from 143 subjects (79 healthy individuals and 64 breast cancer patients) for further analysis. Of the 79 biopsies from healthy women, 5 had been obtained from a benign lesion. By ultrasound and mammography these 5 were described as fibroadenoma (4) or microcalcification (1).

## **RNA-data processing**

The microarrays were scanned by an Agilent scanner (Agilent Technologies, Santa Clara, CA) and processed in Feature Extraction 9.1.3.1 (Agilent Technologies, Santa Clara, CA). Locally weighted scatterplot smoothing (lowess) was used to normalize the data. The normalized and log<sub>2</sub>-transformed data were stored in the Stanford Microarray

Database (SMD)[13] and retrieved from SMD for further statistical analyses. Flagged spots were treated as missing values. The dataset now counted 40791 probes. Clone IDs with 20% or more missing values were excluded. Gene filtering was performed to include only probes with variation across samples, so that probes with less than three arrays being at least 1.6 standard deviations from the mean were excluded. For the 79 healthy women, this probe filtration resulted in an expression dataset of 9767 probes and 79 arrays each representing one individual. For the breast cancer women, a dataset of 64 arrays and 10153 probes were obtained after filtration, and for both groups combined, a dataset of 143 arrays and 13699 probes. Missing values were imputed in R using the method `impute.knn` in the library `impute` [14].

## **Genotyping**

Blood DNA was extracted by phenol/chloroform extraction followed by ethanol precipitation (Nuclear Acid Extractor 340A; Applied Biosystems, Foster City, California, USA) according to standard procedures. *UGT* genotype data was retrieved from two sources: Genome wide association studies (GWAS) using the Human-1 109K BeadChip (Illumina Inc, San Diego, CA) and candidate gene based study using iPlex, Sequenom. For the GWAS, each sample was subject to whole genome amplification using Illumina proprietary reagents [15]. The amplified DNA was fragmented and hybridized according to the protocol. The BeadArray reader (Illumina Inc, San Diego, CA) with the BeadScan software (Illumina Inc, San Diego, CA) was used to image the beadchips. Non-polymorphic probes and probes with more than 20% missing values and were excluded and data processed as described previously [16]. The candidate gene SNP analyses were

performed using the iPLEX assay in conjunction with the Sequenom MassARRAY platform. Multiplexing was performed in 384 plates using 1 ul DNA pr. well with one well containing up to 29 reactions. The technology is described in detail on the sequenom web-page [17].

## **Mammograms**

Routine descriptions of mammograms by local radiologists were collected. Craniocaudal mammograms of both breasts were digitized using a high resolution Kodak Lumisys 85 scanner (Kodak, Rochester, New York). Density was quantified using the University of Southern California Madena assessment method [18]. In brief, the method works as follows: A reader (trained by GU) outlines the total area of the breast using a computerized tool, the software then counts the number of pixels. This represents the total breast area. MD is assessed (by GU), first by identifying a region of interest that incorporates all dense areas except those representing the pectoralis muscle and scanning artifacts, and then by applying a yellow tint to all pixels within the region of interest shaded at or above a threshold intensity of gray. The software then counts the tinted pixels, which represents the area of absolute density. The percent density is the absolute density area divided by the total breast area and is the value used for these analyses. Test-retest reliability was 0.99 for absolute density.

## Statistical analysis

Clustering was performed using MatLab (version R2007b) (The MathWorks Inc., Natick, MA) with Ward linkage and Euclidean distances. Before clustering, the data were gene centred, i.e. for every probe the mean expression across all samples was calculated and was subtracted from the log<sub>2</sub>-ratios for that gene. This was done for visualisation purposes only, clustering with uncentred data returns the same clusters. Significance Analysis of Microarrays (SAM) (version 3.02) [19,20] for Excel was used for analysis of differentially expressed genes between two groups of data. The data were not gene centred for the SAM analysis. A total of 500 permutations were used. Quantitative SAM analysis was used to identify genes differentially expressed according to MD as a continuous variable. Statistical significance tests and regression analysis were performed in R 2.9.0 [21]. To test for difference in the mean of phenotypic variables (MD, age, body mass index (BMI)) in different clusters of women, we used two-sided t-tests (assuming equal variance in the groups) and ANOVA for continuous variables and chi-squared/Fisher's exact tests for categorical variables [22]. To investigate the similarities of distributions of *UGT* genes between tumour samples and normal samples with low MD and high MD respectively, Kullback-Leibler distances between normalized distributions of the histograms of the data were calculated by use of MatLab. The cancer samples in our study were grouped into subtypes and assigned a risk group using the PAM50 gene list published by Parker et al [23]. SNP-analysis was done by use of R 2.9.0. The association between gene expression and SNPs was assessed using expression quantitative trait loci (eQTL) [24] *in cis* (10<sup>6</sup> bp on each side of the gene) using the R package eMap v1.1 [25]. Comparing the akaike information criterion for different models

predicting MD, the lower criterion singled out a linear regression model as the model fitting the distribution of the data best. A linear regression model was fitted in R 2.9.0 with MD as a continuous response variable and the following covariates: UGT2B7, two probes for UGT2B10, UGT2B11, ESR1, age, BMI, current hormone therapy, age at first birth and parity. Gene expression, age, age at first birth and BMI were entered into the model as continuous variables. Stepwise variable selection was performed, starting with all variables included in the model. For every step, the variable with the highest p-value was rejected from the model and the model was refitted. This was repeated until all variables included in the model had a p-value  $<0.05$ . To correct for the influence of age, this variable was forced to stay in the model. A sensitivity analysis was performed excluding extreme ages (30 years or younger) to check the robustness of the data. We also fitted linear regression stratified on age (younger or older than 50 years of age) and current use of hormone therapy. Gene ontology analysis was performed by the use of DAVID Bioinformatics Resources 2008 from the National Institute of Allergy and Infectious Diseases, NIH [26]. Functional annotation clustering was applied and the following gene ontology categories were selected: biological processes (all), molecular function (all) and the KEGG pathway database. We included gene ontology terms with a p-value (FDR-corrected) of  $<0.01$  containing between 5 and 500 genes.

The normalized, log<sub>2</sub>-transformed data are available in Gene Expression Omnibus with accession number [GEO:GSE18672]. The data are not gene centered or gene filtered.

# Results

## Gene expression and mammographic density

To identify genes differentially expressed according to MD we performed quantitative SAM with MD as a continuous variable using gene expression data from the normal biopsies. Of 9767 probes only 25 probes, representing 24 genes, were differentially expressed according to MD, with reduced expression associated with higher MD (false discovery rate (FDR)<25%) (Table 1 and Additional file 1). Gene ontology analysis revealed no significant terms and we found no pathway associated with this gene set. The *UGT* genes and oestrogen receptor gene (*ESR1*) were among the genes significantly down-regulated in breasts with high MD. The percentage of samples with low *UGT* expression was higher in tumour samples than in normal samples with low MD, whereas the percentage was more similar between tumour samples and normal samples with high MD (Additional file 2, Figure S1). The function of *UGT*-enzymes in oestradiol metabolism is illustrated in Figure 1. In healthy women, the expression of the different *UGT* genes was highly correlated with each other and the four probes clustered together (Additional file 2, Figure S2, S3 and Table S1).

MD was lower in women with BMI  $\geq 25$  compared to those with BMI < 25 ( $p=0.01$ ), but unrelated to other epidemiologic variables. *UGT* expression was not significantly associated with age, BMI, age at first birth or current hormone therapy use in the healthy women (Additional file 2, Table S2).

To dissect the impact of age and hormone therapy use, we performed SAM analyses to identify differentially expressed genes according to MD, while stratifying for age and postmenopausal hormone therapy use. For healthy women younger than 50 years of age, the *UGT* genes were not significant at FDR=25%. For healthy women aged 50 or older, 49 probes were significantly down-regulated in breasts with MD of 30% or higher (FDR <25%). Of these, 17 were overlapping with those significantly down-regulated among healthy women in the unstratified analysis. The *UGT* genes were not in this list. We then stratified the women aged 50 or older on current hormone therapy use. When only those currently using hormone therapy were included in the analysis, *UGT2B7* and *UGT2B11* were among the 6 genes differentially expressed with an FDR<10E-5 and *UGT2B28* with FDR <25%. For healthy women above 50 years of age and not currently using hormone therapy, several of the 24 genes were differentially expressed according to MD with an FDR<25%, but again the *UGT* genes were not in this list (Additional file 3).

These analyses were confirmed fitting a linear regression model. While the other variables were excluded from the model with insignificant p-values, age was kept in the model to control for the age-effect. After stepwise variable selection, the only significant variables remaining in the model were *UGT2B10* (A\_23\_P7342)(p=0.005) and BMI (p=0.015). Sensitivity analysis excluding extreme ages (30 years and younger) did not alter the results (*UGT2B10* p=0.003, BMI p=0.016) and indicates the robustness of the results. *ESR1* was borderline significant in both these analyses. These results were not significantly altered when MD was log2-transformed. For further stratification see Table 2.

Unsupervised hierarchical clustering of the 79 samples from healthy women showed two main clusters. MD was not significantly different between these two clusters (Additional file 2, Figure S3).

In the breast cancer group, MD was significantly associated with age and BMI, with higher MD in the younger women and in those with BMI<25. Both MD and UGT expression tended to be higher in women with receptor positive tumours, but this was not significant for any type of receptor. UGT-expression in tumours was unrelated to age, BMI, age at first birth and current hormone therapy (data not shown). There was a higher proportion of oestrogen receptor positive tumours among the breast cancer patients with high MD ( $\geq 30\%$ ) compared with low ( $< 30\%$ ) MD (10/10 vs 36/40, Fisher=0.001). There was no significant association between tumour subtype and level of MD as assessed by ANOVA. There was no indication that degree of MD was associated with the risk of relapse as assessed by the method of Parker et al [23] (Additional file 2, Figure S4).

Nine probes were differentially expressed according to MD in cancer samples (FDR<25%) (Table 3). None of these were overlapping with the 24 genes differentially expressed in the samples from the breasts of healthy women.

## **Genetic polymorphisms**

In order to identify genetic determinants of the expression of the *UGT* genes found to be associated with MD, we performed eQTL analyses of SNPs in these genes as available

from an array based GWAS study and a candidate gene study. 21 SNPs in *UGT* genes were present on the 109K array from Illumina, and 9 SNPs from the candidate gene analysis. Of these, 5 SNPs were associated with the expression of *UGT* genes or other genes in their vicinity at  $p=0.05$  (Additional file 4). Two of these SNPs, both located in *UGT2B10* (rs1828705, rs1828705), were significantly associated with gene expression of another *UGT* gene (*UGT2B7* and *UGT2B28*).

## Discussion

Previously, whole genome expression profiling of normal breast tissue (all cell types included) has been performed to a limited extent [27,28]. Yang et al recently performed a study of cancer-free breast tissue obtained from mastectomies in breast cancer patients with high and low MD [29]. They identified a list of 73 genes differentially expressed between high and low MD samples. Specifically, this included the down-regulation of several TGF $\beta$ -related genes in samples with high MD. In the present study we analysed breast biopsies from 79 healthy women and tumours of 64 women with breast cancer. Twenty-four genes were differentially expressed according to MD in the healthy samples. In breast tumours, none of these 24 genes were found differentially expressed according to MD. Tumour specific deregulation of a large number of mRNA transcripts may be expected to overshadow the MD signature. In addition, the sample size is limited and the two sample sets (cases and controls) are not directly comparable with respect to MD (Additional file 2, Figure S5).

In our study, three *UGT* genes (*UGT2B11*, *UGT2B10* and *UGT2B7*) were differentially expressed according to MD in the breasts of healthy women. All these three enzymes had decreased expression in dense breasts. Previous knowledge links the UGT enzymes to the metabolism of female hormones known to influence the mammary glands (Figure 1). The over-representation of *UGT* genes on the list of significant genes along with a biological link makes these genes particularly interesting. In a linear regression model with age as a confounding factor, BMI and one of two probes for *UGT2B10* were the only significant variables independently predicting MD, with *ESR1* as a borderline significant covariate.

The expression of these three *UGT2B* genes is highly correlated to each other and as expected only one probe remained in the regression model as an independent predictor of MD. BMI is known to be the strongest and most consistent epidemiologic predictor of MD, and is expected to remain in the model. It is noteworthy that one of the *UGT* genes has an independent predictive value of a greater significance and magnitude than BMI. MD is determined by multiple factors. In a study of limited sample size, we can only expect to identify the strongest predictors.

*UGT2B7* is postulated to protect the breast tissue from oestrogen metabolites locally [30], and this is consistent with our findings that breasts with higher MD have reduced expression of this gene. The main metabolites of oestradiol and oestrone (hydroxyl- and methoxy oestrogen compounds) bind to the oestrogen receptor, but with a reduced affinity compared with oestradiol. *UGT2B10* and *11* are not yet reported associated with MD or breast cancer, but *UGT2B10* is involved in the metabolism of tobacco-related nitrosamines [31]. Less is known about and *UGT2B11*. The different *UGT2B* genes are located close to each other on chromosome 4 and there is great homology between the genes (Additional file 2, Figure S6). *UGT1A1*, previously linked to MD and breast cancer [32], is not represented on the microarray used in this study.

We have identified a set of genes differentially expressed according to MD. Interestingly, the *UGT* genes seem, to a greater extent than the other genes, to be more similarly expressed between tumour samples and normal samples from breasts with high MD as compared to normal samples from breasts with low MD (Additional file 2, Table S4 and

Figure S7). The other differentially expressed genes generally express the same levels in the tumours and in the biopsies from the healthy women with low MD. We cannot exclude that the *UGT* genes confer risk for breast cancer development through increasing MD, but further studies would be needed to investigate this.

We found the *UGT* genes to be differentially expressed in young women and women over 50 years of age currently on hormone therapy. SAM analysis of MD in women younger than 50 did not give any differentially expressed genes with an  $FDR < 25\%$ . However, several *UGT*-probes are on the top of the list of genes down-regulated in samples from breasts with high MD. The lack of significance could be due to low sample size ( $n=30$ ). Since *UGT* enzymes conjugate oestradiol metabolites, its effect will be greater when there is an increased level of oestradiol present, whether the oestradiol is endogenous or exogenous. The linear regression analysis showed that *UGT2B10* was predicting MD independent of age in all women, younger women and women older than 50 currently using hormones. This leads to the hypothesis that decreased *UGT* expression in the breast of a woman with increased levels of female hormones confers an increased MD and possibly an increased risk of breast cancer.

The biology in breasts with high and low MD may differ, partly due to differences in proportion of fatty tissue. Therefore, we looked for differentially expressed genes in a subset of samples including only samples from breasts with  $MD > 20\%$ . The fact that the *UGT2B* gene family is so strongly represented among the down-regulated genes (six probes representing five different *UGT2B* genes are the only genes differentially

expressed with an FDR <10E-5) indicate that reduced UGT expression is of greater significance in breasts with higher MD and lower content of fatty tissue.

We find that *ESR1* is down-regulated in biopsies from healthy women with high MD compared to those with low MD. This is not consistent with previous findings [33] and contrary to what one would expect since *ESR1* induces transcription and epithelial growth and high MD may contain increased amounts of epithelial cells [34,35]. However, increased levels of oestradiol have been shown to decrease levels of ESR1 in breast cancer [36], and in normal breast tissue in monkeys [37] and in mice [38]. Increased levels of oestradiol may increase MD. Elevated expression of ESR1 is common postmenopausally [37] and represents non-proliferating cells. The association between reduced levels of ESR1 and high MD may reflect high levels of oestradiol. We found that ESR1 was only a borderline significant predictor of MD in models with stepwise exclusion of covariates. In a model including ESR1 with only age or age and UGT2B10, ESR1 was significantly predicting MD. The independent contribution of ESR1 in predicting MD was significant in older women, where the effect of UGT2B10 was not present. There could be a link between UGT-expression and ESR1-expression in that reduced metabolism of oestradiol-metabolites increase the levels of ESR1-ligands (oestradiol metabolites) and hence reduce ESR1-levels. The UGT-enzyme activity may be the cause of the alterations leading to increased MD by this mechanism. Reduced ESR1 is only borderline significant in predicting MD and could also be an intermediate factor.

MD is the result of complex biological processes without any single determining factor. BMI is the single most important factor found to date, and is also significant in this study. Age seems to have its effect mainly through hormonal influence, except for in postmenopausal women not taking hormones, where age has a significant, independent effect on MD. MD is not significantly different between the two main clusters from unsupervised hierarchical clustering of the samples from healthy women. MD is hence not related to the main variation in the normal samples.

The genes whose expression we have found to be associated with MD do have a fairly high FDR in a SAM analysis and are not significant in all stratified analyses, suggesting that they may play a role in only subsets of individuals and other factors also have a significant contribution. Despite this, in linear regression models UGT2B10 is an independent predictor of MD along with BMI.

There is a substantial heritable proportion of MD. SNPs in *UGT* genes with influence on the UGT expression have been described [8,39]. We identified two UGT-SNPs associated with the expression of other *UGT* genes. Due to their homology and co-localization on the chromosome, they may share common control loci that affect the expression of multiple *UGT* genes. It remains to be investigated in larger and better powered epidemiological studies whether any of these SNPs are associated to MD per se.

We do not know enough about the variability of gene expression within normal breasts to know if the genes relevant for MD are adequately represented by one biopsy taken from

an area with some MD. It is previously shown that two biopsies from the same breast tumour, before and after chemotherapy, cluster together [40]. The tumours may, however, be more homogenous than normal breast tissue. Variability in gene expression within each breast will make it difficult to detect genes with only a minor influence on MD so that only the strongest factors are identified. In an unpublished dataset we found no significant difference between UGT-expression in tumours and normal adjacent tissue tested by paired t-test (Additional file 2, Table S5). This is merely an indication that the expression in one breast might be similar for different locations in the breast and hence be used to look for associations with MD.

In this study, healthy individuals had higher MD than the breast cancer patients. The women recruited in the study had been referred to a breast diagnostic centre for a second look. Since high MD confers an increased risk for breast cancer and mammograms with high MD are more difficult to interpret, they most likely had a higher MD. In addition, the inclusion criterion of some visible MD for biopsy may have influenced the mean MD of the study population. The two populations are not directly comparable with respect to MD and related parameters. This lack of comparability on MD does not affect the analyses of gene expression among the healthy women only.

We obtained good quality microarrays from only 79 of 120 healthy women and from 64 of 66 breast cancer patients. This was due to low mRNA-yield or low mRNA-quality. The biopsies from healthy women consistently yielded less mRNA than the tumour samples. There is significantly higher MD in the breasts of healthy women with

successful microarrays than in those with unsuccessful microarrays (37% vs 29%,  $p=0.03$ ). Since samples from breasts with low MD are under-represented in the microarray study, it is harder to identify genes that are differentially expressed between breast tissue with high and low MD. Despite these limitations, we have identified differentially expressed genes. These genes might have a greater significance than shown in this study.

Normal breast tissue yields less RNA than tumour tissue. The biopsies in this study were small and in agreement with the pathologist, all tissue from normal breasts was prioritized for RNA-extraction rather than histological evaluation. Imprint was not in routine use in the hospitals where we started this study. In order to make it possible for the staff to include women in this study in a busy schedule we had to use procedures already established. We do therefore not have any information about the cell types of the normal biopsies. Knowledge about the cell types present in the biopsies would have facilitated the analysis.

The two UGT2B10-probes behave differently in our dataset. Both probes map to the 3'end of the UGT2B10-gene by BLAT (98.4% homology for A\_23\_P7342 and 100% homology for A\_24\_P521559). The discrepancy in UGT2B10-expression detected by the two probes may be due to the fact that they both also share substantial sequence homology with other, but different UGT2B-genes.

## Conclusions

We have identified a set of genes that are differentially expressed according to MD in breast samples from healthy women. Some of these genes are known to influence MD and breast cancer, such as *ESR1* and *UGT2B7*. Two less described *UGT* genes, *UGT2B10* and *UGT2B11*, are also differentially expressed. The expression of the three *UGT* genes is reduced in samples with high MD and also in tumour samples, but does not vary between different tumour subtypes or risk groups. The *UGT* enzymes are known to conjugate active oestrogen-metabolites. We show that *UGT2B10* expression and BMI are independent predictors of MD. The influence of reduced *UGT* expression was strongest in women under exposure of female hormones. Two candidate SNPs are associated with the *UGT* gene expression *in cis*. We hypothesise that reduced expression of *UGT* genes in women exposed to female sex hormones, increase MD and that this may be associated with an increased risk of breast cancer. Further studies of these genes are needed to test the hypothesis that the gene products from these genes protect the breast from the oestrogen-induced MD and thereby reducing the risk of breast cancer.

## Abbreviations

*ESR1*, oestrogen receptor; FDR, false discovery rate; MD, mammographic density; SAM, significance analysis of microarrays; SNP, single nucleotide polymorphism; *UGT*, uridine 5'-diphospho-glucuronosyltransferase.

## **Competing interests**

The authors declare that they have no competing interests.

## **Authors' contributions**

The trial was designed by ALBD, ÅH, GU, MMH and VNK. ALBD and ÅH ensured funding. MMH, JOF, DN, LR, IKB and VDH assisted in data collection. MB and VNK are responsible for SNP analyses. VDH and TL contributed to the laboratory work. GU estimated the amount of mammographic density. OCL, VDH, HKS and MB have done statistical analyses of the data. ÅH, ALBD and VDH interpreted the results and wrote the paper. All authors were involved in reviewing the report.

## **Acknowledgements**

This study was funded primarily by The Research Council of Norway and South-Eastern Norway Regional Health Authority. We thank all the women who participated in the study and all the personnel in the hospitals who made the inclusion of these women possible, in particular the responsible radiologists: Einar Vigeland, Rolf O Næss and Else Berit Velken. We would also like to thank Lars Ottestad for help in the initiation of the project, Hilde Johnsen, Caroline Jevanord Frøyland and Marit Hilsen for lab assistance and to Eunjung Lee for statistical help. No medical writers were involved in this paper.

## Reference List

1. Oza AM, Boyd NF: **Mammographic parenchymal patterns: a marker of breast cancer risk.** *Epidemiol Rev* 1993, **15**:196-208.
2. Boyd NF, Byng JW, Jong RA, Fishell EK, Little LE, Miller AB, Lockwood GA, Tritchler DL, Yaffe MJ: **Quantitative classification of mammographic densities and breast cancer risk: results from the Canadian National Breast Screening Study.** *J Natl Cancer Inst* 1995, **87**:670-675.
3. Ursin G, Ma H, Wu AH, Bernstein L, Salane M, Parisky YR, Astrahan M, Siozon CC, Pike MC: **Mammographic density and breast cancer in three ethnic groups.** *Cancer Epidemiol Biomarkers Prev* 2003, **12**:332-338.
4. Boyd NF, Lockwood GA, Martin LJ, Knight JA, Byng JW, Yaffe MJ, Tritchler DL: **Mammographic densities and breast cancer risk.** *Breast Dis* 1998, **10**:113-126.
5. Guo YP, Martin LJ, Hanna W, Banerjee D, Miller N, Fishell E, Khokha R, Boyd NF: **Growth factors and stromal matrix proteins associated with mammographic densities.** *Cancer Epidemiol Biomarkers Prev* 2001, **10**:243-248.
6. Ursin G, Lillie EO, Lee E, Cockburn M, Schork NJ, Cozen W, Parisky YR, Hamilton AS, Astrahan MA, Mack T: **The relative importance of genetics and environment on mammographic density.** *Cancer Epidemiol Biomarkers Prev* 2009, **18**:102-112.
7. Guillemette C, Levesque E, Harvey M, Bellemare J, Menard V: **UGT genomic diversity: beyond gene duplication.** *Drug Metab Rev* 2009.
8. Mackenzie PI, Gregory PA, Lewinsky RH, Yasmin SN, Height T, McKinnon RA, Gardner-Stephen DA: **Polymorphic variations in the expression of the chemical detoxifying UDP glucuronosyltransferases.** *Toxicol Appl Pharmacol* 2005, **207**:77-83.
9. Cheng Z, Rios GR, King CD, Coffman BL, Green MD, Mojarrabi B, Mackenzie PI, Tephly TR: **Glucuronidation of catechol estrogens by expressed human UDP-glucuronosyltransferases (UGTs) 1A1, 1A3, and 2B7.** *Toxicol Sci* 1998, **45**:52-57.
10. Yong M, Schwartz SM, Atkinson C, Makar KW, Thomas SS, Newton KM, iello Bowles EJ, Holt VL, Leisenring WM, Lampe JW: **Associations between polymorphisms in glucuronidation and sulfation enzymes and mammographic breast density in premenopausal women in the United States.** *Cancer Epidemiol Biomarkers Prev* 2010, **19**:537-546.

11. Starlard-Davenport A, Lyn-Cook B, Radominska-Pandya A: **Identification of UDP-glucuronosyltransferase 1A10 in non-malignant and malignant human breast tissues.** *Steroids* 2008, **73**:611-620.
12. Wang H, Karesen R, Hervik A, Thoresen SO: **Mammography screening in Norway: results from the first screening round in four counties and cost-effectiveness of a modeled nationwide screening.** *Cancer Causes Control* 2001, **12**:39-45.
13. **Stanford Microarray Database** [<http://genome-www5.stanford.edu/>]
14. **R library impute.knn**  
[<http://rss.acs.unt.edu/Rdoc/library/impute/html/impute.knn.html>]
15. Gunderson KL, Steemers FJ, Lee G, Mendoza LG, Chee MS: **A genome-wide scalable SNP genotyping assay using microarray technology.** *Nat Genet* 2005, **37**:549-554.
16. Nordgard SH, Johansen FE, Alnaes GI, Bucher E, Syvanen AC, Naume B, Borresen-Dale AL, Kristensen VN: **Genome-wide analysis identifies 16q deletion associated with survival, molecular subtypes, mRNA expression, and germline haplotypes in breast cancer patients.** *Genes Chromosomes Cancer* 2008, **47**:680-696.
17. **Sequenom** [<http://www.sequenom.com/>]
18. Ursin G, Astrahan MA, Salane M, Parisky YR, Pearce JG, Daniels JR, Pike MC, Spicer DV: **The detection of changes in mammographic densities.** *Cancer Epidemiol Biomarkers Prev* 1998, **7**:43-47.
19. Tusher VG, Tibshirani R, Chu G: **Significance analysis of microarrays applied to the ionizing radiation response.** *Proc Natl Acad Sci U S A* 2001, **98**:5116-5121.
20. **Significance Analysis of Microarrays** [<http://www-stat.stanford.edu/~tibs/SAM/>]
21. **R project** [r-project.org]
22. Altman DG: *Practical Statistics for Medical Research*. Boca Raton: Chapman & Hall/CRC; 1999.
23. Parker JS, Mullins M, Cheang MC, Leung S, Voduc D, Vickery T, Davies S, Fauron C, He X, Hu Z, Quackenbush JF, Stijleman IJ, Palazzo J, Marron JS, Nobel AB, Mardis E, Nielsen TO, Ellis MJ, Perou CM, Bernard PS: **Supervised risk predictor of breast cancer based on intrinsic subtypes.** *J Clin Oncol* 2009, **27**:1160-1167.

24. Carlborg O, de Koning DJ, Manly KF, Chesler E, Williams RW, Haley CS: **Methodological aspects of the genetic dissection of gene expression.** *Bioinformatics* 2005, **21**:2383-2393.
25. **UNC Biostatistics Software Links**  
[<http://www.bios.unc.edu/~wsun/software.htm>]
26. **DAVID Bioinformatics Resources 6.7** [<http://david.abcc.ncifcrf.gov/>]
27. Sorlie T, Perou CM, Tibshirani R, Aas T, Geisler S, Johnsen H, Hastie T, Eisen MB, van de RM, Jeffrey SS, Thorsen T, Quist H, Matese JC, Brown PO, Botstein D, Eystein Lønning P, Børresen-Dale AL: **Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications.** *Proc Natl Acad Sci U S A* 2001, **98**:10869-10874.
28. Nicolau M, Tibshirani R, Borresen-Dale AL, Jeffrey SS: **Disease-specific genomic analysis: identifying the signature of pathologic biology.** *Bioinformatics* 2007, **23**:957-965.
29. Yang WT, Lewis MT, Hess K, Wong H, Tsimelzon A, Karadag N, Cairo M, Wei C, Meric-Bernstam F, Brown P, Arun B, Hortobagyi GN, Sahin A, Chang JC: **Decreased TGFbeta signaling and increased COX2 expression in high risk women with increased mammographic breast density.** *Breast Cancer Res Treat* 2009.
30. Guillemette C, Belanger A, Lepine J: **Metabolic inactivation of estrogens in breast tissue by UDP-glucuronosyltransferase enzymes: an overview.** *Breast Cancer Res* 2004, **6**:246-254.
31. Kaivosaari S, Toivonen P, Hesse LM, Koskinen M, Court MH, Finel M: **Nicotine glucuronidation and the human UDP-glucuronosyltransferase UGT2B10.** *Mol Pharmacol* 2007, **72**:761-768.
32. Dalhoff K, Buus JK, Enghusen PH: **Cancer and molecular biomarkers of phase 2.** *Methods Enzymol* 2005, **400**:618-627.
33. Verheus M, Maskarinec G, Erber E, Steude JS, Killeen J, Hernandez BY, Cline JM: **Mammographic density and epithelial histopathologic markers.** *BMC Cancer* 2009, **9**:182.
34. Li T, Sun L, Miller N, Nicklee T, Woo J, Hulse-Smith L, Tsao MS, Khokha R, Martin L, Boyd N: **The association of measured breast tissue characteristics with mammographic density and other risk factors for breast cancer.** *Cancer Epidemiol Biomarkers Prev* 2005, **14**:343-349.
35. Rayter Z: **Steroid receptors in breast cancer.** *Br J Surg* 1991, **78**:528-535.

36. Dunbier AK, Anderson H, Ghazoui Z, Folkerd EJ, A'hern R, Crowder RJ, Hoog J, Smith IE, Osin P, Nerurkar A, Parker JS, Perou CM, Ellis MJ, Dowsett M: **Relationship between plasma estradiol levels and estrogen-responsive gene expression in estrogen receptor-positive breast cancer in postmenopausal women.** *J Clin Oncol* 2010, **28**:1161-1167.
37. Cheng G, Li Y, Omoto Y, Wang Y, Berg T, Nord M, Vihko P, Warner M, Piao YS, Gustafsson JA: **Differential regulation of estrogen receptor (ER)alpha and ERbeta in primate mammary gland.** *J Clin Endocrinol Metab* 2005, **90**:435-444.
38. Saji S, Jensen EV, Nilsson S, Rylander T, Warner M, Gustafsson JA: **Estrogen receptors alpha and beta in the rodent mammary gland.** *Proc Natl Acad Sci U S A* 2000, **97**:337-342.
39. Chen G, Dellinger RW, Gallagher CJ, Sun D, Lazarus P: **Identification of a prevalent functional missense polymorphism in the UGT2B10 gene and its association with UGT2B10 inactivation against tobacco-specific nitrosamines.** *Pharmacogenet Genomics* 2008, **18**:181-191.
40. Perou CM, Sorlie T, Eisen MB, van de RM, Jeffrey SS, Rees CA, Pollack JR, Ross DT, Johnsen H, Akslen LA, Fluge O, Pergamenschikov A, Williams C, Zhu SX, Lønning PE, Børresen-Dale AL, Brown PO, Botstein D: **Molecular portraits of human breast tumours.** *Nature* 2000, **406**:747-752.
41. Chen DT, Nasir A, Culhane A, Venkataramu C, Fulp W, Rubio R, Wang T, Agrawal D, McCarthy SM, Gruidl M, Bloom G, Anderson T, White J, Quackenbush J, Yeatman T: **Proliferative genes dominate malignancy-risk gene signature in histologically-normal breast tissue.** *Breast Cancer Res Treat* 2009.
42. Showe MK, Vachani A, Kossenkov AV, Yousef M, Nichols C, Nikonova EV, Chang C, Kucharczuk J, Tran B, Wakeam E, Yie TA, Speicher D, Rom WN, Albelda S, Showe LC: **Gene Expression Profiles in Peripheral Blood Mononuclear Cells Can Distinguish Patients with Non-Small Cell Lung Cancer from Patients with Nonmalignant Lung Disease.** *Cancer Res* 2009.
43. Jones C, Mackay A, Grigoriadis A, Cossu A, Reis-Filho JS, Fulford L, Dexter T, Davies S, Bulmer K, Ford E, Parry S, Budroni M, Palmieri G, Neville AM, O'Hare MJ, Lakhani SR: **Expression profiling of purified normal human luminal and myoepithelial breast cells: identification of novel prognostic markers for breast cancer.** *Cancer Res* 2004, **64**:3037-3045.

**Figure 1: UGTs conjugate oestrogen-substrates into biologically inactive oestrogen glucuronides.** The figure gives a schematic view with focus on glucuronidation and not a complete picture of oestradiol metabolism. Androgens are also inactivated by UGTs, but are left out of this illustration.

**Table 1 Genes differentially expressed according to MD in non-cancer samples**

| Gene symbol  | Agilent ID   | Cytogenetic band |
|--------------|--------------|------------------|
| 729641       | A_24_P932736 | 8p21.1           |
| FLJ10404     | A_23_P427472 | 5q35.3           |
| VPS18        | A_24_P18802  | 15q15.1          |
| UGT2B10      | A_23_P7342   | 4q13.2           |
| CABP7        | A_24_P177236 | 22q12.2          |
| CD86         | A_24_P131589 | 3q13.33          |
| UGT2B11      | A_23_P212968 | 4q13.2           |
| 580687       | A_23_P152570 | 17p11.2          |
| DIAPH2::RPA4 | A_23_P254212 | Xq21.33          |
| LMOD1        | A_32_P199824 | 1q32.1           |
| UGT2B10      | A_24_P521559 | 4q13.2           |
| PIK3R5       | A_23_P66543  | 17p13.1          |
| ATG7         | A_32_P107994 | 3p25.2           |
| LRRC2        | A_23_P155463 | 3p21.31          |
| RBL1         | A_23_P28733  | 20q11.23         |
| NPY1R        | A_23_P69699  | 4q32.2           |
| 810781       | A_23_P144244 | 3q13.33          |
| 593535       | A_32_P80016  | 15q26.1          |
| H2AFJ        | A_23_P204277 | 12p12.3          |
| 666399       | A_32_P35668  | 20p12.3          |
| Transcribed  | A_24_P640617 | 2p25.2           |
| Transcribed  | A_32_P20997  | 20q13.13         |
| UGT2B7       | A_23_P136671 | 4q13             |
| ESR1         | A_23_P309739 | 6q25.1           |
| SAPS1        | A_23_P119448 | 19q13.42         |

MD, mammographic density.

**Table 2: Linear regression analysis of factors predicting MD in all women and stratified for age and hormone therapy use**

| Women in model  | n  | Variables                   | Beta value   | p-value      |
|---|----|-----------------------------|--------------|--------------|
| All women   | 76 | UGT2B10 <sup>1)</sup>       | -0.6         | 0.902        |
|   |    | UGT2B7                      | 1.8          | 0.631        |
|   |    | UGT2B11                     | 4.8          | 0.275        |
|   |    | ESR1                        | -3.8         | 0.055        |
|   |    | <b>UGT2B10<sup>2)</sup></b> | <b>-5.6</b>  | <b>0.005</b> |
|   |    | <b>BMI</b>                  | <b>-1.5</b>  | <b>0.015</b> |
|   |    | age                         | -0.4         | 0.074        |
| 50 years or older                                     | 46 | UGT2B11                     | 0.2          | 0.987        |
|   |    | UGT2B10 <sup>1)</sup>       | 1.0          | 0.946        |
|   |    | UGT2B7                      | 3.5          | 0.486        |
|   |    | UGT2B10 <sup>2)</sup>       | -3.7         | 0.073        |
|   |    | BMI                         | -1.4         | 0.052        |
|   |    | <b>ESR1</b>                 | <b>-6.0</b>  | <b>0.016</b> |
|   |    | age                         | -0.9         | 0.061        |
| 50 years or older,<br>currently on<br>hormone therapy | 11 | UGT2B10 <sup>1)</sup>       | 7.2          | 0.771        |
|   |    | UGT2B11                     | -5.8         | 0.695        |
|   |    | BMI                         | -2.9         | 0.103        |
|   |    | UGT2B7                      | 6.8          | 0.418        |
|   |    | <b>UGT2B10<sup>2)</sup></b> | <b>-27.0</b> | <b>0.000</b> |
|   |    | <b>ESR1</b>                 | <b>-8.1</b>  | <b>0.011</b> |
|   |    | age                         | -0.9         | 0.103        |
| 50 years or older,<br>never used<br>hormone therapy   | 28 | UGT2B11                     | -0.7         | 0.948        |
|   |    | UGT2B10 <sup>1)</sup>       | 3.3          | 0.809        |
|   |    | UGT2B7                      | 3.1          | 0.555        |
|   |    | UGT2B10 <sup>2)</sup>       | -1.4         | 0.607        |
|   |    | BMI                         | -0.9         | 0.348        |
|   |    | <b>ESR1</b>                 | <b>-6.0</b>  | <b>0.033</b> |
|   |    | <b>Age</b>                  | <b>-1.5</b>  | <b>0.004</b> |
| Younger than fifty                                    | 30 | UGT2B7                      | 0.4          | 0.950        |
|   |    | UGT2B10 <sup>1)</sup>       | -1.2         | 0.866        |
|   |    | ESR1                        | -0.9         | 0.835        |
|   |    | UGT2B11                     | 8.4          | 0.225        |
|   |    | BMI                         | -1.4         | 0.216        |

|                             |             |              |
|-----------------------------|-------------|--------------|
| <b>UGT2B10<sup>2)</sup></b> | <b>-6.2</b> | <b>0.040</b> |
| Age                         | -0.3        | 0.610        |

1) A\_24\_P521559, 2) A\_23\_P7342

Factors predicting MD after stepwise exclusion of non-significant factors are shown.

Variables listed in the order of exclusion from the model. P-value from the last equation

including the variable is shown. Age is forced to stay in the model. UGT2B10

(A\_23\_P7342) is a significant, independent predictor of MD in all analysis with a

majority of women under influence of female hormones; women younger than 50 years

of age and women currently on hormone therapy. MD, mammographic density.

**Table 3: Genes differentially expressed according to MD in cancer samples**

| Agilent ID   | Gene name | FDR (%) |
|--------------|-----------|---------|
| A_32_P171923 | 730402    | 0.00    |
| A_32_P480177 | TNN       | 0.00    |
| A_23_P200298 | AGL       | 0.00    |
| A_24_P87036  | TMEM16A   | 0.00    |
| A_23_P312150 | EDN2      | 14.87   |
| A_23_P83388  | EPPK1     | 14.87   |
| A_32_P60065  | F2RL2     | 19.82   |
| A_32_P158272 | MRNA      | 19.82   |
| A_23_P105012 | HRASLS2   | 19.82   |

FDR, false discovery rate; MD, mammographic density.

## **Additional files**

### **Additional file 1**

**Title:** Healthy SAM MD

**Description:** SAM analysis for genes differentially expressed according to MD.

### **Additional file 2**

**Title:** Figures and tables

**Description:** A collection of figures and tables describing the data set and the UGT genes. The main text refers to individual figures and tables in this file.

### **Additional file 3**

**Title:** Healthy SAM MD stratified

**Description:** SAM analysis for gene differentially expressed according to MD stratified on age and use of hormone therapy.

### **Additional file 4**

**Title;** eQTL

**Description:** Expression quantitative trait loci (eQTL) analysis of SNPs affecting the expression of UGT genes *in cis*.




Figure 1

**Additional files provided with this submission:**

Additional file 1: Additional file 1 - Healthy SAM MD.xlsx, 10714K

<http://breast-cancer-research.com/imedia/1764071922403955/supp1.xlsx>

Additional file 2: Additional file 2 - Figures and Tables.pdf, 299K

<http://breast-cancer-research.com/imedia/2581470484410719/supp2.pdf>

Additional file 3: Additional file 3 - Healthy SAM MD stratified.xlsx, 15K

<http://breast-cancer-research.com/imedia/5662988754039577/supp3.xlsx>

Additional file 4: Additional file 4 - eQTL.xlsx, 31K

<http://breast-cancer-research.com/imedia/2058924485403957/supp4.xlsx>