

This Provisional PDF corresponds to the article as it appeared upon acceptance. Copyedited and fully formatted PDF and full text (HTML) versions will be made available soon.

The association of polymorphisms in hormone metabolism pathway genes, menopausal hormone therapy, and breast cancer risk: a nested case-control study in the California Teachers Study cohort

Breast Cancer Research 2011, **13**:R37 doi:10.1186/bcr2859

Eunjung Lee (lee@usc.edu)
Fredrick Schumacher (fschumac@usc.edu)
Juan Pablo Lewinger (lewinger@usc.edu)
Susan L Neuhausen (sneuhausen@coh.org)
Hoda Anton-Culver (hantoncu@uci.edu)
Pamela L Horn-Ross (Pamela.Horn-Ross@cpic.org)
Katherine D Henderson (khenderson@coh.org)
Argyrios Ziogas (aziogas@uci.edu)
David Van Den Berg (dvandenb@usc.edu)
Leslie Bernstein (LBernstein@coh.org)
Giske Ursin (gursin@usc.edu)

ISSN 1465-5411

Article type Research article

Submission date 9 December 2010

Acceptance date 1 April 2011

Publication date 1 April 2011

Article URL <http://breast-cancer-research.com/content/13/2/R37>

This peer-reviewed article was published immediately upon acceptance. It can be downloaded, printed and distributed freely for any purposes (see copyright notice below).

Articles in *Breast Cancer Research* are listed in PubMed and archived at PubMed Central.

For information about publishing your research in *Breast Cancer Research* go to

<http://breast-cancer-research.com/info/instructions/>

© 2011 Lee *et al.*; licensee BioMed Central Ltd.

This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The association of polymorphisms in hormone metabolism pathway genes, menopausal hormone therapy, and breast cancer risk: a nested case-control study in the California Teachers Study cohort

Eunjung Lee¹, Fredrick Schumacher¹, Juan Pablo Lewinger¹, Susan L Neuhausen², Hoda Anton Culver³, Pamela L Horn-Ross⁴, Katherine D Henderson², Argyrios Ziogas³, David Van Den Berg¹, Leslie Bernstein², Giske Ursin^{1,5,6}

¹ Department of Preventive Medicine, Keck School of Medicine, University of Southern California, 1441 Eastlake Ave., Los Angeles, CA 90089, USA

² Department of Population Sciences, Beckman Research Institute, City of Hope, 1500 Duarte Rd, Duarte, CA 91010, USA

³ Department of Epidemiology, School of Medicine, University of California, 224 Irvine Hall, Irvine, CA 92697-7550, USA

⁴ Cancer Prevention Institute of California, 2201 Walnut Ave., Fremont, CA 94538, USA

⁵ Department of Nutrition, University of Oslo, Sognsvannsveien 9, 0372 Oslo, Norway

⁶ Cancer Registry of Norway, PB 5313, Majorstuen, 0304 Oslo, Norway

* Corresponding author: Giske Ursin. E-mail: gursin@usc.edu

Abstract

Introduction: The female sex steroids estrogen and progesterone are important in breast cancer etiology. It therefore seems plausible that variation in genes involved in metabolism of these hormones may affect breast cancer risk, and that these associations may vary depending on menopausal status and use of hormone therapy.

Methods: We conducted a nested case-control study of breast cancer in the California Teachers Study cohort. We analyzed 317 tagging single nucleotide polymorphisms (SNPs) in 24 hormone pathway genes in 2746 non-Hispanic white women: 1351 cases and 1395 controls. Odds ratios (ORs) and 95% confidence intervals (CIs) were estimated by fitting conditional logistic regression models using all women or subgroups of women defined by menopausal status and hormone therapy use. *P* values were adjusted for multiple correlated tests (P_{ACT}).

Results: The strongest associations were observed for SNPs in *SLCO1B1*, a solute carrier organic anion transporter gene, which transports estradiol-17 β -glucuronide and estrone-3-sulfate from the blood into hepatocytes. Ten of 38 tagging SNPs of *SLCO1B1* showed significant associations with postmenopausal breast cancer risk; 5 SNPs (rs11045777, rs11045773, rs16923519, rs4149057, rs11045884) remained statistically significant after adjusting for multiple testing within this gene ($P_{ACT}=0.019-0.046$). In postmenopausal women who were using combined estrogen-progestin therapy (EPT) at cohort enrollment, the OR of breast cancer was 2.31 (95% CI=1.47-3.62) per minor allele of rs4149013 in *SLCO1B1* ($P=0.0003$; within-

gene $P_{ACT}=0.002$; overall $P_{ACT}=0.023$). SNPs in other hormone pathway genes evaluated in this study were not associated with breast cancer risk in premenopausal or postmenopausal women.

Conclusions: We found evidence that genetic variation in *SLCO1B1* is associated with breast cancer risk in postmenopausal women, particularly among those using EPT.

Introduction

Reproductive and hormonal factors, including age at menarche, parity, number of full-term pregnancies, age at first full-term pregnancy, breastfeeding, age at menopause, body mass index (BMI), and physical activity are associated with breast cancer risk [1, 2]. Consistent with these observations, breast cancer risk is higher among women with higher circulating levels of endogenous estrogen [3-5], and among women using combined postmenopausal estrogen and progestin therapy (EPT) [6-11].

Sex steroid hormones, whether endogenous or exogenous, are synthesized and metabolized by many different enzymes (reviewed in [12]). Therefore genetic variation among genes regulating sex steroid hormone levels may increase or decrease breast cancer risk by influencing hormone metabolism. Polymorphisms in several hormone pathway genes, including *CYP19A1* and *COMT*, have been associated with endogenous hormone levels [13-16]; however, studies investigating the association of genetic variation in hormone metabolism genes and breast cancer risk have generated mixed results [14, 17-28]. Recently, a large study from the Breast and Prostate Cancer Cohort Consortium (BPC3) comprehensively analyzed 37 steroid hormone metabolism pathway genes in relation to breast cancer risk and reported null associations [16, 29], suggesting that inconsistencies in the literature may be due to findings observed by chance in small studies. However, it is possible that the inconsistencies may be at least partly explained by differences in the distribution of environmental factors that modify the effects of genetic polymorphisms.

EPT use increases breast cancer risk to a much greater extent than estrogen-only therapy (ET) [6-11]. Therefore, it is important to examine EPT and ET use separately when investigating gene-hormone therapy (HT) interactions. To date, few studies have investigated the gene-HT interactions by hormone formulation using a comprehensive SNP-tagging approach [30]. The California Teachers Study is an effective resource to study these questions because detailed data on hormone use was collected at baseline, and approximately 41% and 28% of the postmenopausal participants reported current use of EPT and ET, respectively [31]. Thus, we systematically investigated whether any of 24 hormone metabolism pathway genes or their interactions with HT was associated with breast cancer risk utilizing data from a case-control study nested within the California Teachers Study.

Materials and methods

Participants

The California Teachers Study has been previously described in detail [32]. Briefly, the California Teachers Study is a prospective cohort of women who were current, recent, or retired California public school professionals in 1995. By returning a baseline questionnaire in 1995-96, 133,479 women joined the cohort and provided detailed information on menopausal status, HT use, and other lifestyle and medical factors. The baseline questionnaire is available on the California Teachers Study website [33]. Cancer diagnoses in the cohort are identified through annual linkage with the California Cancer Registry which identifies at least 99% of cancers diagnosed in California [34]. The California Teachers Study has been approved by the Institutional Review Boards at all participating institutions: the Cancer Prevention Institute of

California (CPIC), the University of California, Irvine (UCI), the University of Southern California (USC), and the City of Hope (COH) in accord with assurances filed with and approved by the U.S. Department of Health and Human Services.

The nested breast cancer case-control study was designed to obtain biospecimens from breast cancer cases and unaffected controls within the 113,590 members of the cohort who were under age 80 years at baseline, had continued residence in California during the study period (1995 to time of blood draw) and, before 1998, had no prior history of invasive or *in situ* breast cancer. Cases were women diagnosed with a histologically confirmed invasive primary carcinoma of the breast (International Classification of Disease for Oncology code C50 restricted to morphology codes under 8590) at age 80 years or younger, between January 1, 1998 and May 31, 2007. One control participant per case was randomly selected from the cohort and frequency-matched to the case on age at baseline (within 5 year age groups), self-reported race/ethnicity (white, African American, Latina, Asian, other), and three broad geographic region (i.e. California Teachers Study specimen collection centers). Cancer cases were identified through quarterly linkages of the cohort to the California Cancer Registry database. Control selection was conducted on a quarterly basis without replacement. For each wave of control selection, a reference date was determined (i.e., January, April, July and October of each year). Approximately equal numbers of controls were selected in each wave. One control participant was diagnosed with breast cancer after her control selection and was included in the analysis both as a control and a case.

Collection of biological specimens and DNA extraction

Collection of biological specimens was conducted at three study centers (CPIC in the northern half of California, USC and UCI in the southern half). Women who declined blood draw were asked whether they were willing to provide a saliva sample and if so, an Oragene DNA self-collection kit (DNA Genotek, Ontario, Canada) was mailed to the participant with informed consent and return postage paid mailing materials. From the 8,118 eligible participants (2,618 cases and 5,500 selected controls), we collected biological specimens for 74% of the cases (1923 cases; 1684 blood specimens and 239 saliva specimen) and 61% of the controls (3350 controls; 3012 blood specimens and 338 saliva specimens). All biologic samples were sent via overnight courier to the UCI laboratory for DNA extraction. DNA was extracted from blood clots using Qiagen Clotspin Baskets and DNA QIAmp DNA Blood maxi kits (Qiagen Inc) following Qiagen's protocols. DNA was extracted from saliva samples using the Oragene protocol (DNA Genotek, Ontario, Canada). The nested breast cancer case-control study of the California Teachers Study has been approved by the Institutional Review Boards at all participating institutions, and all participants provided written informed consent.

Tagging SNP selection and genotyping

We investigated 24 genes that are involved in female sex steroid hormone biosynthesis, metabolism, or excretion. Reviews on the function of these genes are available elsewhere [12, 35, 36]. For 21 of these genes, a tagging SNP approach was used (see Supplementary Table S1 in Additional file 1). For 16 of the 21 genes, we selected linkage disequilibrium tagging SNPs across each gene, 20 kb upstream of 5' of the UTR (untranslated region) and 10 kb downstream of 3' of the UTR. The tagging SNPs were selected to capture all common SNPs (minor allele frequency $\geq 5\%$) in individuals of European ancestry with minimum pairwise $R^2 \geq 0.80$ using the

Snagger software [37] and the data from the International HapMap Project for white CEPH (Utah residents with ancestry from northern and western Europe) population (Hapmap, release 21, July 2006, genotype build 36 [38]). Tagging SNPs for five genes included in the current study had been selected by BPC3 using the TagSNPs program [29, 39]. To facilitate the comparison across studies, we used the BPC3-selected tagging SNPs for these five genes (see Supplementary Table S1 in Additional file 1). The BPC3-selected tagging SNPs captured all common SNPs ($MAF \geq 5\%$) in whites with minimum pairwise $R^2 \geq 0.8$. For the remaining three genes of the 24 genes, we genotyped a few selected potentially functional SNPs because these were lower priority and we had a finite number of SNPs available on our platform, or for CYP19A1, because it had been thoroughly analyzed [13] (see Supplementary Table S1 in Additional file 1).

A total of 1751 breast cancer cases and 1697 controls were available for genotyping. We included a random sample of 193 replicates (105 cases and 88 controls) to monitor reproducibility and track plate flips or switches. The DNA samples were genotyped for the selected tagging SNPs using the Illumina Golden Gate Assay (Illumina, Inc., San Diego, CA) in the USC Core Facility. About 10% of the genotyped samples, including 189 cases and 150 controls, had a genotyping success rate (call rate) less than 90% and were excluded from the analyses. The genotyping concordance rate based on the 160 duplicate samples with call rate $\geq 90\%$ was 99.9%. Call rates were lower when the DNA was obtained from saliva samples: 23% of saliva samples and 8% of blood samples had a call rate less than 90% and these samples were therefore excluded from the analyses. However, the genotyping concordance after excluding the low call rate samples was excellent for saliva samples ($>99.9\%$). In addition, results from

sensitivity analyses excluding saliva samples were similar to those using all samples. For the current study, we also excluded 88 women (52 cases and 36 controls) who self-reported to have had a previous history of cancer, leaving 1,510 cases and 1,511 controls. Because the majority (~91%) of participants was non-Hispanic whites, we restricted the analyses to 2,746 non-Hispanic white women (1,351 cases and 1,395 controls).

Of the 355 SNPs genotyped, 332 SNPs had a SNP-call rate $\geq 90\%$. We excluded an additional three SNPs that had discordant readings in more than 2 duplicate pairs, eight SNPs with MAF $< 1\%$ among non-Hispanic white controls, and four SNPs in *COMT*, *CYP11A*, *SULT1A1*; *SULT1A2*, *UGT1A8* not in Hardy Weinberg equilibrium (HWE $P < 0.001$), thus analyzing 317 SNPs in the current study.

For 19 of the 20 genes for which we used the tagging SNP approach, the genotyped tagging SNPs efficiently captured 70-100% of all common SNPs (MAF $> 5\%$) in the Hapmap dataset of European ancestry (Hapmap release 24, genotype build 36) with pairwise $R^2 \geq 0.80$ (Supplementary Table S1 in Additional file 1). We did not have sufficient tagging coverage for *CYP21A2*.

Imputation

We imputed SNPs in gene regions where we found a statistically significant association ($P < 0.01$ before multiple testing correction) with breast cancer risk among all women or among subgroups defined by menopausal status. To do this, we used publicly available Hapmap genotype data in the CEPH population as the reference sample (Hapmap release 24, genotype build 36 [38]) and

MACH 1.0 [40]. We excluded imputed SNPs when the minor allele frequency (MAF) was < 1% or when the R^2 was < 0.30 [40].

Statistical analyses

We used conditional logistic regression models with strata defined by 5-year age group and the three specimen collection center to estimate the odds ratios (ORs), 95% confidence intervals (95% CIs) and P-values associated with each SNP using log-additive models. The results did not change after further adjustment for potential confounders including menopausal status (premenopausal, postmenopausal, unknown), HT use status at baseline (never used HT, currently using ET, currently using EPT, used HT in the past, unknown), BMI (<25, 25-<30, ≥ 30 kg/m², unknown), parity (0, 1~2, ≥ 3 , unknown), and oral contraceptive use (never, ever, unknown). Therefore we presented the results from the conditional logistic regression models not adjusting for these potential confounders.

We performed subgroup-specific analyses by menopausal status, and among postmenopausal women, by HT use at baseline, defining the groups of interest as: never used HT, currently using ET, and currently using EPT. We calculated P for interaction by likelihood ratio test comparing the model with and without the product term of genotype (0, 1, and 2 copies of minor allele, as ordinal variable) and menopausal status or HT use. The interaction tests for HT use were done separately for current EPT use (as compared to never HT use) and for current ET use (as compared to never HT use).

Results

A greater proportion of breast cancer cases than controls were currently using EPT at baseline; a lower proportion of cases than controls had high parity. Cases also had slightly earlier age at menarche than control women (Table 1).

Evaluation of the q-q plot of the P-values for the association between the 317 SNPs in the 24 hormone metabolism genes and breast cancer risk showed no evidence of systematic bias (Supplementary Figure S1 in Additional file 2).

We observed statistically significant associations at $P < 0.01$ with 2 SNPs in *SLCO1B1* and 1 SNP in *HSD17B4* in the overall analysis (Table 2). However, after correcting for multiple comparisons, none of these associations was statistically significant. For postmenopausal women, 10 SNPs in *SLCO1B1* were associated with breast cancer risk with an uncorrected $P < 0.01$. Of these, the associations for SNPs rs11045773, rs11045777, rs16923519, rs4149057, rs11045884 remained statistically significant after correction for multiple testing within the gene (within-gene $P_{ACT} < 0.05$). However, after correcting for multiple testing across all genes, none of these associations was statistically significant. The ORs and 95% CIs associated for all tested SNPs are provided in Supplementary Table S2 in Additional file 3. Of these, there was some evidence that rs4149013 in *SLCO1B1* was associated with breast cancer risk in postmenopausal women (OR=1.39, 95% CI=1.07-1.81, uncorrected $P=0.015$).

When examined by HT use, we observed a strong association between several SNPs in *SLCO1B1* and postmenopausal breast cancer risk in current EPT users (Table 3). Breast cancer

risk among postmenopausal women who were using EPT at baseline increased more than two-fold per minor allele of rs4149013 (OR=2.31, 95% CI=1.47-3.62, P = 0.0003, within-gene P_{ACT} = 0.002). This association was statistically significant even after P_{ACT} adjustment across all SNPs studied (P_{ACT} = 0.024). The P value for interaction (EPT versus never HT use) was 0.019 (not corrected for multiple testing). When we combined the homozygous and heterozygous minor allele carriers (i.e. a dominant genetic model), we observed similar OR estimates and P values (OR=2.43, 95% CI=1.53-3.85, P=0.0002; Supplementary Table S3 in Additional file 4). We did not observe any significant associations in never HT users and ET users. There was no statistically significant difference in effects when stratifying by estrogen receptor status.

Discussion

In this case-control study nested within the California Teachers Study cohort, genetic variation in only one (*SLCO1B1*) of 24 genes in the hormone metabolism pathway genes was associated with breast cancer risk. *SLCO1B1*, a gene involved in the hepatic uptake of female sex steroids, seemed to be associated with breast cancer risk among postmenopausal women. This association was statistically significant after correcting for multiple testing within the gene but was not statistically significant after we corrected for multiple testing across genes. However, there was also an indication that combined HT may interact with SNPs in *SLCO1B1*; one variant in *SLCO1B1* (rs4149013) was statistically significantly associated with breast cancer risk in EPT users.

Our findings of no association between SNPs in hormone metabolism pathway genes and breast cancer risk are consistent with the results from other large studies such as BPC3 [13, 29, 30, 41] and meta-analyses of selected functional SNPs in *CYP1A1* [23], *SULT1A1* [42, 43], *CYP1B1* [44, 45], and *COMT* [46], although two smaller meta-analyses of selected functional SNP in *COMT* supported an association in Caucasian populations [47, 48]. While a few studies have suggested associations between genetic polymorphisms in other hormone pathway genes, including *CYP11A* [25, 26], *CYP1A1/CYP1A2* [49, 50], *CYP1B1* [51, 52], *SULT1E1* [53], or *COMT* [54], many of these associations were not observed consistently [29, 49, 50, 55-57]. In addition, the few studies other than BPC3 that have investigated polymorphisms in *CYP2C9* [51], *CYP3A4* [49, 51, 52], *HSD17B2* [58], *SRD5A1* [52], and *UGT2B7* [51], in relation to breast cancer risk among Caucasian populations, have reported no associations. However, a recent study using admixture maximum likelihood (AML)-based global tests reported that genetic variation in androgen-estrogen conversion pathway was associated with breast cancer risk, although no single SNP was significant after correcting multiple testing [59].

To our knowledge no studies have investigated genetic variation in *SLCO1B1* and breast cancer risk by hormone therapy use. *SLCO1B1*, also known as *OATP-C* or *SLC21A6*, is expressed in the liver, and plays an important role in transporting drugs and endogenous substrates from the blood into the hepatocytes (reviewed in [60]). Endogenous substrates of *SLCO1B1* include steroid hormone conjugates such as estradiol-17 β -glucuronide and estrone-3-sulfate [61, 62]. Serum estrone sulfate (E1S) is a major form of circulating estrogen in postmenopausal women and can be converted to estradiol in breast tissue [63]. E1S is also a major component of conjugated equine estrogens, the estrogen component of the predominant (prior to 2002) HT

regimens in the US [64]. Genetic variation in *SLCO1B1* has been shown to decrease the uptake of E1S and estradiol glucuronide in several [61, 65] but not all [66] studies. Further, one study has shown that genetic variation in *SLCO1B1* is associated with blood E1S levels in Caucasians [61], suggesting that genetic variation in *SLCO1B1* may interact with HT use. Rs4149013 is located near the 5' end of *SLCO1B1*. The functional significance of this variant is not known, but even if there is none, this variant could be linked to a causal allele.

In the publicly-available Cancer Genetic Markers of Susceptibility (CGEMS) breast cancer data [67], we found additional support implicating *SLCO1B1*. In CGEMS, five genotyped SNPs in *SLCO1B1* (rs704166, rs852550, rs852549, rs7489119, rs2306283) were associated with breast cancer risk with $P < 0.05$. Although these 5 SNPs, as imputed genotypes, were null in our dataset (data not shown), this could be due to misclassification from imputation or false-positive associations across both CGEMS and our data. However, our findings and those of CGEMS, combined with the previous literature on the role of this gene in affecting estrone absorption, suggest that further investigation of the role of *SLCO1B1* genetic variation and its interaction with EPT on breast cancer risk is warranted.

The strengths of this study include the systematic investigation of a large number of hormone metabolism genes and the detailed information on HT use collected at baseline. A limitation of our study was the inability to genotype all tagging SNPS for several of the genes of interest, including *AKR1C4*, *ARSC*, and *CYP19A1*. Thus, we cannot exclude the possibility that the lack of associations for these loci were due to incomplete tagging. Overall, we had 80% statistical power to detect ORs ranging from 1.17 to 1.40 for SNPs with MAF of 0.05 to 0.49 using log-additive models and an alpha of 0.05. For the subset analyses among postmenopausal women,

the minimum detectable OR ranged from 1.20 to 1.46 for SNPs with $MAF \geq 0.05$. The statistical power to detect associations in premenopausal women or to detect interactions with menopausal status or HT use was limited. Another limitation of this study is that HT use status assessed at baseline may have changed during follow-up. The participation rate (donating biological specimen for this nested case-control study) among the potentially eligible cohort members was moderate (74% for cases; 61% for controls). However, it is unlikely that the participation was differential according to genotype and case status, and thus selection bias is unlikely to have influenced our findings.

Conclusions

Common genetic variations in *SLCO1B1* may be associated with breast cancer risk in postmenopausal women, particularly in EPT users. The known effects of variants in *SLCO1B1* on estrogen metabolism suggest that further study of the role of *SLCO1B1* is warranted.

Abbreviations

OR, odds ratio; CI, confidence interval; P_{ACT} , P values adjusted for multiple correlated tests; BMI, body mass index; EPT, estrogen progestin combined therapy; ET, estrogen-therapy; HT, hormone therapy; CPIC, Cancer Prevention Institute of California; UCI, University of California, Irvine; USC, University of Southern California; COH, City of Hope National Medical Center; SNP, single nucleotide polymorphism; MAF, minor allele frequency; HWE, Hardy-Weinberg equilibrium; BPC3, Breast and Prostate Cancer Cohort Consortium.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

EL performed the statistical analysis and drafted the manuscript. FS, JPL, and SN participated in the statistical analysis and drafted the manuscript. PLHR, HAC, and LB participated in the design of the nested case-control study and were responsible for specimen collection and obtained grant funding for the study. KDH prepared analytic variables including menopausal status and hormone therapy use. AZ participated in coordinating specimen process and SNP selection. DVDB coordinated tagging SNP selection and performed the genotyping. GU participated in the design of the study, SNP selection, and supervised the analysis and manuscript preparation. All authors edited the manuscript and provided comments on the intellectual content, and have approved the final manuscript.

Acknowledgements

This research was supported by grant R01 CA77398 and CA17054, and K05 CA 136967 from the National Cancer Institute and contract 97-10500 from the California Breast Cancer Research Fund. EL was supported by California Breast Cancer Research Program 14FB-0104. The funding sources did not contribute to the design or conduct of the study, nor to the writing or submission of this manuscript.

The collection of cancer incidence data used in this study was supported by the California Department of Health Services as part of the statewide cancer reporting program mandated by California Health and Safety Code Section 103885; the National Cancer Institute's Surveillance, Epidemiology and End Results Program under contract N01-PC-35136 awarded to the Cancer

Prevention Institute of California (formerly the Northern California Cancer Center), contract N01-PC-35139 awarded to the University of Southern California, and contract N02-PC-15105 awarded to the Public Health Institute; and the Centers for Disease Control and Prevention's National Program of Cancer Registries, under agreement #U55/CCR921930-02 awarded to the Public Health Institute. The ideas and opinions expressed herein are those of the authors and endorsement by the California Health and Human Services Agency, the National Cancer Institute, and the Centers for Disease Control and Prevention or their contractors and subcontractors is not intended nor should be inferred.

The authors would like to thank the California Teachers Study Steering Committee who are responsible for the formation and maintenance of the cohort within which this study was conducted but who did not directly contribute to the current paper: Ellen T. Chang, Christina A. Clarke, Dennis Deapen, James V. Lacey Jr, Huiyan Ma, David O. Nelson, Rich Pinder, Peggy Reynolds, Daniel O. Stram, Dee W. West, and Sophia S. Wang.

References

1. Collaborative Group on Hormonal Factors in Breast Cancer: **Breast cancer and breastfeeding: collaborative reanalysis of individual data from 47 epidemiological studies in 30 countries, including 50302 women with breast cancer and 96973 women without the disease.** *Lancet* 2002, **360**:187-195.
2. Kelsey JL, Gammon MD, John EM: **Reproductive factors and breast cancer.** *Epidemiologic Reviews* 1993, **15**:36-47.
3. Key T, Appleby P, Barnes I, Reeves G: **Endogenous sex hormones and breast cancer in postmenopausal women: reanalysis of nine prospective studies.** *J Natl Cancer Inst* 2002, **94**:606-616.
4. Tworoger SS, Missmer SA, Barbieri RL, Willett WC, Colditz GA, Hankinson SE: **Plasma sex hormone concentrations and subsequent risk of breast cancer among women using postmenopausal hormones.** *J Natl Cancer Inst* 2005, **97**:595-602.
5. Eliassen AH, Missmer SA, Tworoger SS, Spiegelman D, Barbieri RL, Dowsett M, Hankinson SE: **Endogenous steroid hormone concentrations and risk of breast cancer among premenopausal women.** *J Natl Cancer Inst* 2006, **98**:1406-1415.
6. **Breast cancer and hormone replacement therapy: collaborative reanalysis of data from 51 epidemiological studies of 52,705 women with breast cancer and 108,411 women without breast cancer. Collaborative Group on Hormonal Factors in Breast Cancer.** *Lancet* 1997, **350**:1047-1059.
7. Ross RK, Paganini-Hill A, Wan PC, Pike MC: **Effect of hormone replacement therapy on breast cancer risk: estrogen versus estrogen plus progestin.** *J Natl Cancer Inst* 2000, **92**:328-332.
8. Schairer C, Lubin J, Troisi R, Sturgeon S, Brinton L, Hoover R: **Menopausal estrogen and estrogen-progestin replacement therapy and breast cancer risk.** *Jama* 2000, **283**:485-491.
9. Olsson HL, Ingvar C, Bladstrom A: **Hormone replacement therapy containing progestins and given continuously increases breast carcinoma risk in Sweden.** *Cancer* 2003, **97**:1387-1392.
10. Farquhar CM, Marjoribanks J, Lethaby A, Lamberts Q, Suckling JA: **Long term hormone therapy for perimenopausal and postmenopausal women.** *Cochrane Database Syst Rev* 2005:CD004143.
11. Rossouw JE, Anderson GL, Prentice RL, LaCroix AZ, Kooperberg C, Stefanick ML, Jackson RD, Beresford SA, Howard BV, Johnson KC, Kotchen JM, Ockene J: **Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results from the Women's Health Initiative randomized controlled trial.** *JAMA* 2002, **288**:321-333.
12. Zhu BT, Conney AH: **Functional role of estrogen metabolism in target cells: review and perspectives.** *Carcinogenesis* 1998, **19**:1-27.
13. Haiman CA, Dossus L, Setiawan VW, Stram DO, Dunning AM, Thomas G, Thun MJ, Albanes D, Altshuler D, Ardanaz E, Boeing H, Buring J, Burt N, Calle EE, Chanock S, Clavel-Chapelon F, Colditz GA, Cox DG, Feigelson HS, Hankinson SE, Hayes RB, Henderson BE, Hirschhorn JN, Hoover R, Hunter DJ, Kaaks R, Kolonel LN, Le Marchand L, Lenner P, Lund E, et al.: **Genetic variation at the CYP19A1 locus predicts circulating estrogen levels but not breast cancer risk in postmenopausal women.** *Cancer Res* 2007, **67**:1893-1897.

14. Dunning AM, Dowsett M, Healey CS, Tee L, Luben RN, Folkard E, Novik KL, Kelemen L, Ogata S, Pharoah PD, Easton DF, Day NE, Ponder BA: **Polymorphisms associated with circulating sex hormone levels in postmenopausal women.** *J Natl Cancer Inst* 2004, **96**:936-945.
15. Tworoger SS, Chubak J, Aiello EJ, Ulrich CM, Atkinson C, Potter JD, Yasui Y, Stapleton PL, Lampe JW, Farin FM, Stanczyk FZ, McTiernan A: **Association of CYP17, CYP19, CYP1B1, and COMT polymorphisms with serum and urinary sex hormone concentrations in postmenopausal women.** *Cancer Epidemiol Biomarkers Prev* 2004, **13**:94-101.
16. Beckmann L, Husing A, Setiawan VW, Amiano P, Clavel-Chapelon F, Chanock SJ, Cox DG, Diver R, Dossus L, Feigelson HS, Haiman C, Hallmans G, Hayes RB, Henderson BE, Hoover RN, Hunter DJ, Khaw K, Kolonel LN, Kraft P, Lund E, Le Marchand L, Peeters PH, Riboli E, Stram D, Thomas G, Thun MJ, Tumino R, Trichopoulos D, Vogel U, Willett WC, et al.: **Comprehensive analysis of hormone and genetic variation in 36 genes related to steroid hormone metabolism in pre- and postmenopausal women from the breast and prostate cancer cohort consortium (BPC3).** *J Clin Endocrinol Metab* 2011, **96**:E360-367.
17. Dunning AM, Healey CS, Pharoah PD, Teare MD, Ponder BA, Easton DF: **A systematic review of genetic polymorphisms and breast cancer risk.** *Cancer Epidemiol Biomarkers Prev* 1999, **8**:843-854.
18. Haiman CA, Hankinson SE, Spiegelman D, Brown M, Hunter DJ: **No association between a single nucleotide polymorphism in CYP19 and breast cancer risk.** *Cancer Epidemiol Biomarkers Prev* 2002, **11**:215-216.
19. Haiman CA, Hankinson SE, Spiegelman D, Colditz GA, Willett WC, Speizer FE, Kelsey KT, Hunter DJ: **The relationship between a polymorphism in CYP17 with plasma hormone levels and breast cancer.** *Cancer Res* 1999, **59**:1015-1020.
20. Becchis M, Frairia R, Ferrera P, Fazzari A, Ondei S, Alfarano A, Coluccia C, Biglia N, Sismondi P, Fortunati N: **The additionally glycosylated variant of human sex hormone-binding globulin (SHBG) is linked to estrogen-dependence of breast cancer.** *Breast Cancer Res Treat* 1999, **54**:101-107.
21. Forsti A, Jin Q, Grzybowska E, Soderberg M, Zientek H, Sieminska M, Rogozinska-Szczepka J, Chmielik E, Utracka-Hutka B, Hemminki K: **Sex hormone-binding globulin polymorphisms in familial and sporadic breast cancer.** *Carcinogenesis* 2002, **23**:1315-1320.
22. Low YL, Dunning AM, Dowsett M, Luben RN, Khaw KT, Wareham NJ, Bingham SA: **Implications of gene-environment interaction in studies of gene variants in breast cancer: an example of dietary isoflavones and the D356N polymorphism in the sex hormone-binding globulin gene.** *Cancer Res* 2006, **66**:8980-8983.
23. Masson LF, Sharp L, Cotton SC, Little J: **Cytochrome P-450 1A1 gene polymorphisms and risk of breast cancer: a HuGE review.** *Am J Epidemiol* 2005, **161**:901-915.
24. Haiman CA, Stram DO, Pike MC, Kolonel LN, Burt NP, Altshuler D, Hirschhorn J, Henderson BE: **A comprehensive haplotype analysis of CYP19 and breast cancer risk: the Multiethnic Cohort.** *Hum Mol Genet* 2003, **12**:2679-2692.
25. Setiawan VW, Cheng I, Stram DO, Giorgi E, Pike MC, Van Den Berg D, Pooler L, Burt NP, Le Marchand L, Altshuler D, Hirschhorn J, Henderson BE, Haiman CA: **A systematic assessment of common genetic variation in CYP11A and risk of breast cancer.** *Cancer Res* 2006, **66**:12019-12025.
26. Zheng W, Gao YT, Shu XO, Wen W, Cai Q, Dai Q, Smith JR: **Population-based case-control study of CYP11A gene polymorphism and breast cancer risk.** *Cancer Epidemiol Biomarkers Prev* 2004, **13**:709-714.
27. Kristensen VN, Borresen-Dale AL: **Molecular epidemiology of breast cancer: genetic variation in steroid hormone metabolism.** *Mutat Res* 2000, **462**:323-333.

28. Pharoah PD, Tyrer J, Dunning AM, Easton DF, Ponder BA: **Association between common variation in 120 candidate genes and breast cancer risk.** *PLoS Genet* 2007, **3**:e42.
29. Canzian F, Cox DG, Setiawan VW, Stram DO, Ziegler RG, Dossus L, Beckmann L, Blanche H, Barricarte A, Berg CD, Bingham S, Buring J, Buys SS, Calle EE, Chanock SJ, Clavel-Chapelon F, Delancey JO, Diver WR, Dorransoro M, Haiman CA, Hallmans G, Hankinson SE, Hunter DJ, Husing A, Isaacs C, Khaw KT, Kolonel LN, Kraft P, Le Marchand L, Lund E, et al.: **Comprehensive analysis of common genetic variation in 61 genes related to steroid hormone and insulin-like growth factor-I metabolism and breast cancer risk in the NCI breast and prostate cancer cohort consortium.** *Hum Mol Genet* 2010, **19**:3873-3884.
30. Feigelson HS, Cox DG, Cann HM, Wacholder S, Kaaks R, Henderson BE, Albanes D, Altshuler D, Berglund G, Berrino F, Bingham S, Buring JE, Burt NP, Calle EE, Chanock SJ, Clavel-Chapelon F, Colditz G, Diver WR, Freedman ML, Haiman CA, Hankinson SE, Hayes RB, Hirschhorn JN, Hunter D, Kolonel LN, Kraft P, LeMarchand L, Linseisen J, Modi W, Navarro C, et al.: **Haplotype analysis of the HSD17B1 gene and risk of breast cancer: a comprehensive approach to multicenter analyses of prospective cohort studies.** *Cancer Res* 2006, **66**:2468-2475.
31. Saxena T, Lee E, Henderson KD, Clarke CA, West D, Marshall SF, Deapen D, Bernstein L, Ursin G: **Menopausal hormone therapy and subsequent risk of specific invasive breast cancer subtypes in the California Teachers Study.** *Cancer Epidemiol Biomarkers Prev* 2010, **19**:2366-2378.
32. Bernstein L, Allen M, Anton-Culver H, Deapen D, Horn-Ross PL, Peel D, Pinder R, Reynolds P, Sullivan-Halley J, West D, Wright W, Ziogas A, Ross RK: **High breast cancer incidence rates among California teachers: results from the California Teachers Study (United States).** *Cancer causes & control* 2002, **13**:625-635.
33. **California Teachers Study.** [<http://calteachersstudy.org/>]
34. Kwong S, Perkin C, Morris C, Cohen R, Allen M, Wright W: **Cancer in California: 1988-1999.** Sacramento (CA): California Department of Health Services, Cancer Surveillance Section; 2001.
35. Li J, Eriksson L, Humphreys K, Czene K, Liu J, Tamimi RM, Lindström S, Hunter DJ, Vachon CM, Couch FJ, Scott CG, Lagiou P, Hall P: **Genetic variation in the estrogen metabolic pathway and mammographic density as an intermediate phenotype of breast cancer.** *Breast Cancer Res* 2010, **12**:R19.
36. Payne AH, Hales DB: **Overview of steroidogenic enzymes in the pathway from cholesterol to active steroid hormones.** *Endocr Rev* 2004, **25**:947-970.
37. Edlund CK, Lee WH, Li D, Van Den Berg DJ, Conti DV: **Snagger: a user-friendly program for incorporating additional information for tagSNP selection.** *BMC Bioinformatics* 2008, **9**:174.
38. **International Hapmap Project.** [<http://hapmap.ncbi.nlm.nih.gov>]
39. Stram DO, Haiman CA, Hirschhorn JN, Altshuler D, Kolonel LN, Henderson BE, Pike MC: **Choosing haplotype-tagging SNPs based on unphased genotype data using a preliminary sample of unrelated subjects with an example from the Multiethnic Cohort Study.** *Hum Hered* 2003, **55**:27-36.
40. Li Y, Abecasis GR: **Mach 1.0: Rapid Haplotype Reconstruction and Missing Genotype Inference.** *Am J Hum Genet* 2006, **S79**:2290.
41. Setiawan VW, Schumacher FR, Haiman CA, Stram DO, Albanes D, Altshuler D, Berglund G, Buring J, Calle EE, Clavel-Chapelon F, Cox DG, Gaziano JM, Hankinson SE, Hayes RB, Henderson BE, Hirschhorn J, Hoover R, Hunter DJ, Kaaks R, Kolonel LN, Kraft P, Ma J, Le Marchand L, Linseisen J, Lund E, Navarro C, Overvad K, Palli D, Peeters PH, Pike MC, et al.: **CYP17 genetic variation and risk of breast and prostate cancer from the National Cancer Institute Breast and Prostate Cancer Cohort Consortium (BPC3).** *Cancer Epidemiol Biomarkers Prev* 2007, **16**:2237-2246.

42. Sun Y, Zang Z, Xu X, Zhang Z, Zhong L, Zan W, Zhao Y, Sun L: **The association of SULT1A1 codon 213 polymorphism and breast cancer susceptibility: meta-analysis from 16 studies involving 23,445 subjects.** *Breast Cancer Res Treat* 2010.
43. Wang Z, Fu Y, Tang C, Lu S, Chu WM: **SULT1A1 R213H polymorphism and breast cancer risk: a meta-analysis based on 8,454 cases and 11,800 controls.** *Breast Cancer Res Treat* 2009.
44. Economopoulos KP, Sergentanis TN: **Three polymorphisms in cytochrome P450 1B1 (CYP1B1) gene and breast cancer risk: a meta-analysis.** *Breast Cancer Res Treat* 2010.
45. Yao L, Fang F, Wu Q, Zhong Y, Yu L: **No association between CYP1B1 Val432Leu polymorphism and breast cancer risk: a meta-analysis involving 40,303 subjects.** *Breast Cancer Res Treat* 2009.
46. Mao C, Wang XW, Qiu LX, Liao RY, Ding H, Chen Q: **Lack of association between catechol-O-methyltransferase Val108/158Met polymorphism and breast cancer risk: a meta-analysis of 25,627 cases and 34,222 controls.** *Breast Cancer Res Treat* 2010, **121**:719-725.
47. Onay UV, Aaltonen K, Briollais L, Knight JA, Pabalan N, Kilpivaara O, Andrulis IL, Blomqvist C, Nevanlinna H, Ozcelik H: **Combined effect of CCND1 and COMT polymorphisms and increased breast cancer risk.** *BMC Cancer* 2008, **8**:6.
48. Ding H, Fu Y, Chen W, Wang Z: **COMT Val158Met polymorphism and breast cancer risk: evidence from 26 case-control studies.** *Breast Cancer Res Treat* 2010.
49. Le Marchand L, Donlon T, Kolonel LN, Henderson BE, Wilkens LR: **Estrogen metabolism-related genes and breast cancer risk: the multiethnic cohort study.** *Cancer Epidemiol Biomarkers Prev* 2005, **14**:1998-2003.
50. Rebbeck TR, Troxel AB, Walker AH, Panossian S, Gallagher S, Shatalova EG, Blanchard R, Norman S, Bunin G, DeMichele A, Berlin M, Schinnar R, Berlin JA, Strom BL: **Pairwise combinations of estrogen metabolism genotypes in postmenopausal breast cancer etiology.** *Cancer Epidemiol Biomarkers Prev* 2007, **16**:444-450.
51. The MARIE-GENICA Consortium on Genetic Susceptibility for Menopausal Hormone Therapy Related Breast Cancer Risk: **Genetic polymorphisms in phase I and phase II enzymes and breast cancer risk associated with menopausal hormone therapy in postmenopausal women.** *Breast Cancer Res Treat* 2010, **119**:463-474.
52. Reding KW, Weiss NS, Chen C, Li CI, Carlson CS, Wilkerson HW, Farin FM, Thummel KE, Daling JR, Malone KE: **Genetic polymorphisms in the catechol estrogen metabolism pathway and breast cancer risk.** *Cancer Epidemiol Biomarkers Prev* 2009, **18**:1461-1467.
53. Choi JY, Lee KM, Park SK, Noh DY, Ahn SH, Chung HW, Han W, Kim JS, Shin SG, Jang IJ, Yoo KY, Hirvonen A, Kang D: **Genetic polymorphisms of SULT1A1 and SULT1E1 and the risk and survival of breast cancer.** *Cancer Epidemiol Biomarkers Prev* 2005, **14**:1090-1095.
54. Ji Y, Olson J, Zhang J, Hildebrandt M, Wang L, Ingle J, Fredericksen Z, Sellers T, Miller W, Dixon JM, Brauch H, Eichelbaum M, Justenhoven C, Hamann U, Ko Y, Bruning T, Chang-Claude J, Wang-Gohrke S, Schaid D, Weinshilboum R: **Breast cancer risk reduction and membrane-bound catechol O-methyltransferase genetic polymorphisms.** *Cancer Res* 2008, **68**:5997-6005.
55. Justenhoven C, Hamann U, Schubert F, Zapatka M, Pierl CB, Rabstein S, Selinski S, Mueller T, Ickstadt K, Gilbert M, Ko YD, Baisch C, Pesch B, Harth V, Bolt HM, Vollmert C, Illig T, Eils R, Dippon J, Brauch H: **Breast cancer: a candidate gene approach across the estrogen metabolic pathway.** *Breast Cancer Res Treat* 2008, **108**:137-149.
56. Reding KW, Li CI, Weiss NS, Chen C, Carlson CS, Duggan D, Thummel KE, Daling JR, Malone KE: **Genetic variation in the progesterone receptor and metabolism pathways and hormone therapy in relation to breast cancer risk.** *Am J Epidemiol* 2009, **170**:1241-1249.

57. Gaudet MM, Chanock S, Lissowska J, Berndt SI, Peplonska B, Brinton LA, Welch R, Yeager M, Bardin-Mikolajczak A, Garcia-Closas M: **Comprehensive assessment of genetic variation of catechol-O-methyltransferase and breast cancer risk.** *Cancer Res* 2006, **66**:9781-9785.
58. Jansson A, Carlsson J, Olsson A, Storm P, Margolin S, Gunnarsson C, Stenmark-Askmalin M, Lindblom A, Persson B, Stal O: **A new polymorphism in the coding region of exon four in HSD17B2 in relation to risk of sporadic and hereditary breast cancer.** *Breast Cancer Res Treat* 2007, **106**:57-64.
59. Low YL, Li Y, Humphreys K, Thalamuthu A, Darabi H, Wedren S, Bonnard C, Czene K, Iles MM, Heikkinen T, Aittomaki K, Blomqvist C, Nevanlinna H, Hall P, Liu ET, Liu J: **Multi-variant pathway association analysis reveals the importance of genetic determinants of estrogen metabolism in breast and endometrial cancer susceptibility.** *PLoS Genet* 2010, **6**:e1001012.
60. Kalliokoski A, Niemi M: **Impact of OATP transporters on pharmacokinetics.** *Br J Pharmacol* 2009, **158**:693-705.
61. van der Deure WM, Friesema EC, de Jong FJ, de Rijke YB, de Jong FH, Uitterlinden AG, Breteler MM, Peeters RP, Visser TJ: **Organic anion transporter 1B1: an important factor in hepatic thyroid hormone and estrogen transport and metabolism.** *Endocrinology* 2008, **149**:4695-4701.
62. Tamai I, Nozawa T, Koshida M, Nezu J, Sai Y, Tsuji A: **Functional characterization of human organic anion transporting polypeptide B (OATP-B) in comparison with liver-specific OATP-C.** *Pharm Res* 2001, **18**:1262-1269.
63. Gamage N, Barnett A, Hempel N, Duggleby RG, Windmill KF, Martin JL, McManus ME: **Human sulfotransferases and their role in chemical metabolism.** *Toxicol Sci* 2006, **90**:5-22.
64. Speroff L, Fritz MA: *Clinical gynecologic endocrinology and infertility.* 7 edn. Philadelphia: Lippincott Williams & Wilkins; 2005.
65. Tirona RG, Leake BF, Merino G, Kim RB: **Polymorphisms in OATP-C: identification of multiple allelic variants associated with altered transport activity among European- and African-Americans.** *J Biol Chem* 2001, **276**:35669-35675.
66. Nozawa T, Nakajima M, Tamai I, Noda K, Nezu J, Sai Y, Tsuji A, Yokoi T: **Genetic polymorphisms of human organic anion transporters OATP-C (SLC21A6) and OATP-B (SLC21A9): allele frequencies in the Japanese population and functional analysis.** *J Pharmacol Exp Ther* 2002, **302**:804-813.
67. Hunter DJ, Kraft P, Jacobs KB, Cox DG, Yeager M, Hankinson SE, Wacholder S, Wang Z, Welch R, Hutchinson A, Wang J, Yu K, Chatterjee N, Orr N, Willett WC, Colditz GA, Ziegler RG, Berg CD, Buys SS, McCarty CA, Feigelson HS, Calle EE, Thun MJ, Hayes RB, Tucker M, Gerhard DS, Fraumeni JF, Jr., Hoover RN, Thomas G, Chanock SJ: **A genome-wide association study identifies alleles in FGFR2 associated with risk of sporadic postmenopausal breast cancer.** *Nat Genet* 2007, **39**:870-874.
68. Conneely KN, Boehnke M: **So Many Correlated Tests, So Little Time! Rapid Adjustment of P Values for Multiple Correlated Tests.** *Am J Hum Genet* 2007, **81**:1158-1168.

Table 1. Characteristics of the study participants at time of joining the cohort

Characteristics	Control	(%)	Case	(%)
N	1,395		1,351	
Age (mean±SD)	56.1 ± 9.5		55.0 ± 9.4	
Menopausal status				
Premenopausal	347	25.3	364	27.5
Postmenopausal*	1024	74.7	962	72.5
Unknown	24	-	25	-
HT use, among postmenopausal women				
Never used HT	153	15.8	112	12.4
Current ET use	277	28.7	230	25.5
Current EPT use	401	41.5	468	51.9
Former ET or EPT use	122	12.6	80	8.9
Ever used Progestin alone	13	1.3	12	1.3
Unknown	58	-	60	-
Parity / total number of full-term pregnancies (FTP)				
Nulligravid	221	16.1	219	16.4
Gravid, nulliparous	61	4.4	66	4.9
1 FTP	159	11.6	174	13.0
2 FTP	487	35.4	472	35.3
3 FTP	274	20.0	277	20.7
4+ FTP	172	12.5	128	9.6
Unknown	21	-	15	-
BMI (kg/m ²)				
<25	765	54.8	778	58.9
25-<30	387	27.7	384	29.1
30+	195	14.0	159	12.0
Unknown	48	-	30	-
Age at menarche (yrs)				
≤10	100	7.3	100	7.5
11-12	600	43.6	599	44.8
13-14	565	41.0	542	40.6
15-16	102	7.4	82	6.1
17+	10	0.7	13	1.0
Unknown/Never had menarche	18	-	15	-
Family history of breast cancer (first-degree relative)				
No	1,167	86.1	1,085	82.1
Yes	189	13.9	236	17.9
Unknown	39	-	30	-
History of breast biopsy				
No	1,086	77.8	1,034	76.5
Yes	309	22.2	317	23.5
Screening mammograms within last 2 years				
No	138	10.0	146	10.9

Yes	1241	90.0	1193	89.1
Unknown	16	-	12	-

* Includes peri-menopausal women.

Abbreviation: FTP, full-term pregnancies

Table 2. Odds ratios (OR) and 95% confidence intervals (CI) for SNPs that were associated with breast cancer risk with P-values < 0.01 (uncorrected for multiple testing), in either overall analyses or in subgroup analyses of premenopausal or postmenopausal women*

Locus	All subjects						Premenopausal						Postmenopausal						
	WW ca/co	VW ca/co	OR (95%CI)	P _{ACT} [†]	P	P _{ACT} [‡]	WW ca/co	VW ca/co	OR (95%CI)	P	P _{ACT} [†]	WW ca/co	VW ca/co	OR (95%CI)	P	P _{ACT} [‡]			
SLCO1B1																			
rs11045773	898/857	401/468	50/67	0.83 (0.72-0.94)	0.004	0.088	227/206	118/127	18/13	0.94 (0.72-1.21)	0.61	1.00	659/636	273/332	29/54	0.77 (0.65-0.90)	0.000	0.021	0.17
rs11045777	899/855	401/471	50/64	0.83 (0.73-0.95)	0.005	0.099	227/204	118/129	18/11	0.94 (0.73-1.22)	0.66	1.00	660/636	273/333	29/53	0.77 (0.65-0.90)	0.000	0.01	0.17
rs976754	1,126/1,207	217/183	8/5	1.27 (1.03-1.55)	0.01	0.26	309/294	50/51	5/2	1.04 (0.71-1.53)	0.76	1.00	795/894	164/127	3/3	1.40 (1.10-1.79)	0.004	0.07	0.15
rs16923519	836/921	452/421	62/53	1.17 (1.02-1.33)	0.01	0.26	244/222	107/114	12/11	0.92 (0.70-1.20)	0.53	1.00	577/685	335/297	50/42	1.26 (1.08-1.47)	0.001	0.03	0.029
rs4149057	496/569	641/638	209/186	1.14 (1.02-1.27)	0.02	0.28	152/133	164/172	48/42	0.96 (0.77-1.19)	0.70	1.00	334/429	466/453	157/140	1.22 (1.07-1.39)	0.002	0.04	0.047
rs4149058	778/854	468/467	95/72	1.15 (1.02-1.30)	0.02	0.34	220/204	122/120	21/22	0.94 (0.73-1.20)	0.66	1.00	541/636	342/337	71/50	1.23 (1.07-1.43)	0.004	0.07	0.052
rs11045884	880/871	412/452	41/63	0.86 (0.75-0.99)	0.03	0.36	216/220	126/107	15/16	1.09 (0.84-1.42)	0.42	1.00	649/638	277/335	25/46	0.78 (0.66-0.92)	0.002	0.04	0.033
rs11045825	953/943	357/394	38/55	0.87 (0.75-1.00)	0.04	0.45	237/236	112/97	14/13	1.09 (0.83-1.42)	0.54	1.00	700/692	237/288	23/42	0.79 (0.66-0.93)	0.004	0.06	0.037
rs10841767	889/874	411/451	49/67	0.88 (0.77-1.00)	0.04	0.47	221/222	125/107	17/18	1.07 (0.83-1.38)	0.51	1.00	653/639	277/334	31/48	0.80 (0.69-0.94)	0.005	0.09	0.058
rs11045813	917/918	372/401	37/56	0.88 (0.76-1.01)	0.06	0.58	225/229	118/99	13/12	1.12 (0.85-1.47)	0.40	1.00	677/676	245/293	23/44	0.79 (0.67-0.94)	0.005	0.08	0.024
HSD17B4																			
rs382719	393/457	672/685	275/241	1.16 (1.04-1.29)	0.008	0.097	105/124	179/172	76/51	1.32 (1.06-1.64)	0.012	0.12	282/329	479/499	194/184	1.11 (0.98-1.26)	0.1	0.65	0.16
rs17388769	919/1,002	393/364	33/21	1.21 (1.04-1.41)	0.01	0.13	243/258	108/89	11/0	1.56 (1.15-2.13)	0.004	0.051	656/729	280/266	22/21	1.12 (0.94-1.34)	0.1	0.7	0.074

* Based on conditional logistic regression models stratified by age group (within 5-yr age groups) and specimen collection centers (CPIC, USC, UCI), using log-additive genetic models.

[†] P_{ACT} (P values adjusted for correlated tests) within each gene was calculated using the methods by Conneely and Boehnke [68]. When applying P_{ACT} across all genes evaluated, none were statistically significant.

Abbreviation: SNP, single nucleotide polymorphism

Table 3. Odds ratios (OR) and 95% confidence intervals (CI) for SNPs that were associated with breast cancer risk with P-values < 0.01 (uncorrected for multiple testing) in subgroup analyses of postmenopausal women who were using EPT at baseline*

Gene	SNP	Never used HT			Using ET			Using EPT			
		N** (ww/vv/vv)	OR (95% CI)	P‡	N** (ww/vv/vv)	OR (95% CI)	P‡	N** (ww/vv/vv)	OR (95% CI)	P‡ (P _{ACT} [¶] ; P _{ACT} [§])	P _{int} [†]
SLCO1B1	rs4149013	114/12/1 148/20/1	0.79 (0.39-1.60)	0.51	199/31/0 245/31/0	1.20 (0.69-2.03)	0.54	393/71/1 373/27/1	2.31 (1.47-3.62)	0.0003 (0.002; 0.023)	0.019
SLCO1B1	rs976754	110/16/1 143/25/1	0.87 (0.45-1.67)	0.67	198/32/0 246/31/0	1.21 (0.71-2.05)	0.48	375/91/2 351/49/1	1.71 (1.19-2.47)	0.004 (0.045; 0.37)	0.11
SLCO1B1	rs11045777	86/35/6 104/56/8	0.68 (0.42-1.09)	0.11	155/66/9 180/82/14	0.88 (0.64-1.19)	0.40	327/131/10 248/134/19	0.72 (0.56-0.92)	0.009 (0.10; 0.68)	0.95
SLCO1B1	rs11045773	86/35/6 103/56/9	0.66 (0.41-1.06)	0.09	155/66/9 181/81/14	0.88 (0.65-1.20)	0.42	326/131/10 248/134/19	0.72 (0.56-0.92)	0.009 (0.11; 0.70)	0.86

* Based on conditional logistic regression models stratified by age group (within 5-yr age groups) and specimen collection centers (CPIC, USC, UCI), using log-additive genetic models.

** N of cases carrying 0, 1, and 2 copies of minor allele and N of controls carrying 0, 1, and 2 copies of minor allele, respectively.

‡ P values not corrected for multiple testing

¶ P_{ACT} (P values adjusted for correlated tests) was calculated within each gene using the methods by Conneely and Boehnke [68].

§ P_{ACT} (P values adjusted for correlated tests) was calculated across all genes using the methods by Conneely and Boehnke [68].

† P values for interaction with never HT users.

Abbreviations: HT, hormone therapy; ET, estrogen-only therapy; EPT, estrogen-progestin therapy; P_{ACT}, P values adjusted for correlated tests

Additional files

Additional file 1

Title: Supplementary Table S1.

Description: A word document of Supplementary Table S1.

Additional file 2

Title: Supplementary Figure S1.

Description: A word document of Supplementary Figure S1.

Additional file 3

Title: Supplementary Table S2.

Description: An excel document of Supplementary Table S2.

Additional file 4

Title: Supplementary Table S3.

Description: A word document of Supplementary Table S3.

Additional files provided with this submission:

Additional file 1: Format_revision_SupplementaryTable1.docx, 19K
<http://breast-cancer-research.com/imedia/1674497824535773/supp1.docx>

Additional file 2: Format_revision_SupplementaryFig1.docx, 29K
<http://breast-cancer-research.com/imedia/4128497835357731/supp2.docx>

Additional file 3: Supplementary table 2_simple_version.xlsx, 61K
<http://breast-cancer-research.com/imedia/138683335357732/supp3.xlsx>

Additional file 4: Format_revision_Supplementary Table 3.doc, 36K
<http://breast-cancer-research.com/imedia/1417198419535772/supp4.doc>